

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> <b>C07K 5/08, 5/10, 7/06, A61K 38/06, 38/07, 38/08</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 95/15973</b> <b>(43) International Publication Date:</b> 15 June 1995 (15.06.95)
<b>(21) International Application Number:</b> PCT/US94/13943 <b>(22) International Filing Date:</b> 5 December 1994 (05.12.94)  <b>(30) Priority Data:</b> 08/164,101                      6 December 1993 (06.12.93)      US Not furnished                      2 December 1994 (02.12.94)      US  <b>(71) Applicant:</b> CYTEL CORPORATION [US/US]; 3525 John Hopkins Court, San Diego, CA 92121 (US).  <b>(72) Inventors:</b> ARRHENIUS, Thomas, S.; 3909 San Augustine Way, San Diego, CA 92130 (US). ELICES, Mariano, J.; 11596 Prairie Shadow Point, San Diego, CA 92126 (US). GAETA, Federico, C., A.; 3226 Rim Rock Circle, Olivenhain, CA 92024 (US).  <b>(74) Agent:</b> CAMPBELL, Cathryn; Campbell and Flores, Suite 700, 4370 La Jolla Village Drive, San Diego, CA 92122 (US).		<b>(81) Designated States:</b> AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> CS-1 PEPTIDOMIMETICS, COMPOSITIONS AND METHODS OF USING THE SAME  <b>(57) Abstract</b>  The present invention contemplates a peptide that inhibits the binding between the VLA-4 receptor expressed on inflammatory leukocytes and the fibronectin CS-1 peptide expressed on endothelial cells that are involved in immunoinflammatory disease states. Pharmaceutical compositions containing a contemplated peptide and processes for treating immunoinflammatory conditions using a binding-inhibitory peptide are also disclosed.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

- 1 -

CS-1 PEPTIDOMIMETICS, COMPOSITIONS  
AND METHODS OF USING THE SAME

5

Description

Technical Field

10           The present invention relates to binding of  
inflammatory cells to endothelial cells that express the  
CS-1 portion of fibronectin on their surfaces, and more  
particularly to the inhibition of that binding by  
peptidomimetic compounds of minimal length.

15

Background Art

          The immune response relies on leukocyte  
trafficking and immune surveillance as one of the  
underpinnings of host defense. Not only does this  
20   immune surveillance allow leukocytes to recirculate  
through lymphoid tissues normally, but also permits  
rapid leukocyte recruitment and extravasation to  
adjacent tissues at sites of inflammation. The  $\alpha 4 \beta 1$   
(CD49d/CD29, VLA-4) cell adhesion receptor is an active  
25   participant in these leukocyte trafficking functions  
[Hemler, Ann. Rev. Immunol., 8:365-400 (1990); Hemler et  
al., Immunol. Rev., 114:45-65 (1990)].

          The VLA-4 integrin heterodimer was discovered  
independently by three research groups and identified as  
30   a surface antigen on lymphocytes [Sanchez-Madrid et al.,  
Eur. J. Immunol., 16:1343-1349 (1986); Clayberger et  
al., J. Immunol., 138:1510-1514 (1987); Hemler et al.,  
J. Biol. Chem., 262:11478-11485 (1987)]. Within the  
integrin family, VLA-4 is unique on several counts: (i)

- 2 -

in contrast to related members of the  $\beta 1$  subfamily, VLA-4 is predominantly expressed on cells of the hematopoietic lineage [Hemler, Ann. Rev. Immunol., 8:365-400 (1990)], and is functionally involved in cell-cell, as well as cell-extracellular matrix (ECM) adhesive interactions [Hemler, Ann. Rev. Immunol., 8:365-400 (1990)]; (ii) despite sequence homology with other integrin  $\alpha$  subunits, the  $\alpha 4$  subunit stands apart from the two major structural clusters of  $\alpha$  subunits because  $\alpha 4$  lacks an inserted I-domain, and does not undergo post-translational cleavage near the transmembrane region [Hemler, Ann. Rev. Immunol., 8:365-400 (1990); Hynes, Cell, 69:11-25 (1992)]; and (iii)  $\alpha 4$  contains a trypsin-like cleavage site that results in cell type-specific surface expression of at least two different structural variants termed  $\alpha 4$ -150 and  $\alpha 4$ -80/70 [Pulido et al., FEBS Lett., 294:121-124 (1991); Teixido et al., J. Biol. Chem., 267:1786-1791 (1992); Rubio et al., Eur. J. Immunol., 22:1099-1102 (1992)].

The VLA-4 integrin appears to be one of the earliest adhesion receptors found on CD34-expressing hematopoietic stem cells [Teixido et al., J. Clin. Invest., 90:358-367 (1992)]. However, VLA-4 is expressed only on mature T and B lymphocytes, natural killer (NK) cells, monocytes, basophils and eosinophils, but not on erythrocytes, platelets and neutrophils [Hemler, Ann. Rev. Immunol., 8:365-400 (1990); Gismondi et al., J. Immunol., 146:384-392 (1991); Walsh et al., J. Immunol., 146:3419-3423 (1991); Bochner et al., J. Exp. Med., 173:1553-1556 (1992); Dobrina et al., J. Clin. Invest., 88:20-26 (1991); Weller et al., Proc. Natl. Acad. Sci. USA, 88:7430-7433 (1991)].

To date, most adhesion functions mediated by VLA-4 can be explained by a direct molecular interaction between the VLA-4 integrin and either of two separate

- 3 -

counterreceptor structures, namely, the cytokine-inducible vascular cell adhesion molecule-1 (VCAM-1) [Elices et al., Cell, 60:577-584 (1990); Rice et al., J. Exp. Med., 171:1369-1374 (1990); Schwartz et al., J. Clin. Invest., 85:2019-2022 (1990); Carlos et al., Blood, 76:965-970 (1990)], and a subset of the ubiquitous ECM protein fibronectin [Wayner et al., J. Cell Biol., 109:1321-1330 (1989); Guan et al., Cell, 60:53-61 (1990); Ferreira et al., J. Exp. Med., 171:351-356 (1990); Elices et al., Cell, 60:577-584 (1990)].

VCAM-1 is a member of the immunoglobulin (Ig) gene superfamily [Osborn et al., Cell, 59:1203-1211 (1989); Rice et al., Science, 246:1303-1306 (1989)] that is expressed predominantly in vascular endothelium in response to pro-inflammatory cytokines such as IL-1, TNF $\alpha$ , and IL-4 [Osborn et al., Cell, 59:1203-1211 (1989); Rice et al., Science, 246:1303-1306 (1989); Thornhill et al., J. Immunol., 145:865-872 (1990); Masinovsky et al., J. Immunol., 145:2886-2895 (1990); Thornhill et al., J. Immunol., 146:592-598 (1991); Schleimer et al., J. Immunol., 148:1086-1092 (1992); Birdsall et al., J. Immunol., 148:2717-2723 (1992); Swerlick et al., J. Immunol., 149:798-705 (1992); Briscoe et al., J. Immunol., 149:2954-2960 (1992)]. The VLA-4 binding sites on VCAM-1 have been mapped to the outermost N-terminal (first) Ig-like region of the 6-Ig-like domain VCAM-1 isoform [Taichman et al., Cell Regul., 2:347-355 (1991); Vonderheide et al., J. Exp. Med., 175:1433-1442 (1992); Osborn et al., J. Exp. Med., 176:99-107 (1992)], and the first and fourth N-terminal Ig-like regions of the 7-Ig-like domain VCAM-1 isoform [Vonderheide et al., J. Exp. Med., 175:1433-1442 (1992); Osborn et al., J. Exp. Med., 176:99-107 (1992)]. Discrete amino acid sequences within the two separate

- 4 -

Ig-like domains in VCAM-1 recognized by the VLA-4 integrin remain to be defined.

In contrast, a high affinity peptide recognition sequence for VLA-4 within fibronectin (FN) has been identified [Wayner et al., J. Cell. Biol., 109:1321-1330 (1989); Ferreira et al., J. Exp. Med., 171:351-356 (1990); Guan et al., Cell, 60:53-61 (1990); Mould et al., J. Biol. Chem., 265:4020-4024 (1990); Garcia-Pardo et al., J. Immunol., 144:3361-3366 (1990); Komoriya et al., J. Biol. Chem., 266:15075-15079 (1991)]. That sequence comprises a 25-amino acid residue stretch, termed CS-1 [Humphries et al., J. Cell Biol., 103:2637-2647 (1986); Humphries et al., J. Biol. Chem., 262:6886-6892 (1987)].

The FN gene contains three separate exons termed EIIIA, EIIIB and V or IIICS, which are subject to alternative splicing [Hynes, "Fibronectin", Springer-Verlag, New York (1990)]. The presence of additional acceptor and donor splice signals within the IIICS region permits generation of increased diversity in FN by virtue of multiple IIICS polypeptide variants, for instance, five in human FN [Vibe-Pedersen et al., FEBS Lett., 207:287-291 (1987); Hershberger et al., Mol. Cell. Biol., 10:662-671 (1990)]. Consequently, only a subset of these molecular variants expresses the 25-amino acid CS-1 sequence recognized by VLA-4 [Wayner et al., J. Cell. Biol., 109:1321-1330 (1989); Guan et al., Cell, 60:53-61 (1990)].

A minimal essential sequence for specific VLA-4 recognition of CS-1 has been identified as the tripeptide Leu-Asp-Val (LDV) [Komoriya et al., J. Biol. Chem., 266:15075-15079 (1991); Wayner et al., J. Cell. Biol., 116:489-497 (1992); Wayner WO 91/03252 published March 21, 1991; Wayner WO 93/12809 published July 8, 1993; and Humphries WO 92/13887, published August 20,

- 5 -

1992] albeit VLA-4 binds to LDV with at least two orders of magnitude lower affinity than to the native CS-1 25-mer. Nowlin et al., J. Biol. Chem., 268(1):20352-20359 (1993) recently described a cystine-linked cyclic pentapeptide said to inhibit binding by both the Arg-Gly-Asp and CS-1 regions of fibronectin.

VLA-4 shares with other members of the  $\beta 1$  integrin subfamily the ability to promote binding and penetration of microbial pathogens into mammalian cells. Thus, specific interactions of  $\beta 1$  integrins with the bacterial protein invasins [Isberg et al., Cell, 60:861-871 (1990); Ennis et al., J. Exp. Med., 177:207-212 (1993)], as well as the protozoan Trypanosoma cruzi [Fernandez et al., Eur. J. Immunol., 23:552-557 (1993)] have been described.

A multitude of in vitro studies suggest interactions of VLA-4 with its two known ligands, VCAM-1 and CS-1 FN, have profound biological significance. For instance, VLA-4 binding to VCAM-1 has been demonstrated in adhesion to cytokine-stimulated vascular endothelium by lymphocytes [Elices et al., Cell, 60:577-584 (1990); Rice et al., J. Exp. Med., 171:1369-1374 (1990); Schwartz et al., J. Clin. Invest., 85:2019-2022 (1990); Carlos et al., Blood, 76:965-970 (1990); Shimizu et al., J. Cell Biol., 113:1203-1212 (1991)], monocytes [Carlos et al., Blood, 77:2266-2271 (1991); Jonjic et al., J. Immunol., 148:2080-2083 (1992)], natural killer (NK) cells [Allavena et al., J. Exp. Med., 173:439-448 (1991)], and eosinophils [Walsh et al., J. Immunol., 146:3419-3423 (1991); Bochner et al., J. Exp. Med., 173:1553-1556 (1992); Dobrina et al., J. Clin. Invest., 88:20-26 (1991); Weller et al., Proc. Natl. Acad. Sci. USA, 88:7430-7433 (1991)]. Because of its involvement in mediating leukocyte-endothelial attachment,

- 6 -

VLA-4/VCAM-1 interactions are considered key in inflammation.

The VLA-4/CS-1 interaction, in turn, has been widely documented in hematopoiesis where adhesive interactions between hematopoietic progenitors expressing VLA-4 [Hemler et al., Immunol. Rev., 114:45-65 (1990); Williams et al., Nature, 352:438-441 (1991); Roldan et al., J. Exp. Med., 175:1739-1747 (1992); Sawada et al., J. Immunol., 149:3517-3524 (1992); Wadsworth et al., J. Immunol., 150:847-857 (1993)] and their ECM microenvironment play a critical role in precursor maturation and differentiation. Thus, CS-1 peptides have been shown to inhibit (i) attachment of murine hematopoietic stem cells to ECM derived from bone marrow stroma [Williams et al., Nature, 352:438-441 (1991)], (ii) immunoglobulin secretion by bone marrow-derived B cell progenitors [Roldan et al., J. Exp. Med., 175:1739-1747 (1992)], (iii) bursal and postbursal development of chicken B cells [Palojoki et al., Eur. J. Immunol., 23:721-726 (1993)], and (iv) thymocyte adhesion and differentiation induced by thymic stromal cell monolayers [Utsumi et al., Proc. Natl. Acad. Sci. USA, 88:5685-5689 (1991); Sawada et al., J. Immunol., 149:3517-3524 (1992)]. VLA-4/CS-1 may also be involved in embryonic development, because CS-1 peptides have been shown to interfere with migration of avian neural crest cells [Dufour et al., EMBO J., 7:2661-2671 (1988)].

In addition to VCAM-1, FN and CS-1 have also been implicated in the pathology of rheumatoid arthritis (RA) [Laffon et al., J. Clin. Invest., 88:546-552 (1992)]. A role for the CS-1 splicing variant of FN has been established in mediating migration of inflammatory cells such as eosinophils across endothelial cell



- 7 -

monolayers of VLA-4-expressing leukocytes [Kuijpers et al., J. Exp. Med., 178:279-284 (1993)].

5 The vast body of work suggesting that VLA-4 plays a role in leukocyte trafficking and inflammation has been largely confirmed by in vivo studies using anti-VLA-4 antibodies in various animal models. Essentially, the skin, brain, kidney, lung and gut are targets of a wide variety of VLA-4-dependent inflammatory reactions mostly resulting from recruitment  
10 of mononuclear leukocytes and eosinophils.

More specifically, these in vivo studies are as follows: contact hypersensitivity (CH) and delayed type hypersensitivity (DTH) in the mouse and rat [Ferguson et al., Proc. Natl. Acad. Sci. USA, 88:8072-  
15 8076 (1991); Issekutz, Cell Immunol., 138:300-312 (1991); Issekutz, J. Immunol., 147:4178-4184 (1991); Elices et al., Clin. Exp. Rheumatol., 11:S77-80 (1993); Chisholm, et al., Eur. J. Immunol., 23:682-688 (1993)]; experimental autoimmune encephalomyelitis (EAE) in the  
20 mouse and rat [Yednock et al., Nature, 356:63-66 (1992); Baron et al., J. Exp. Med., 177:57-68 (1993)]; nephrotoxic nephritis in the rat [Mulligan et al., J. Clin. Invest., 91:577-587 (1993)]; passive cutaneous anaphylaxis in the guinea pig [Weg et al., J. Exp. Med.,  
25 177:561-566 (1993)]; immune complex-induced lung injury in the rat [Mulligan et al., J. Immunol., 150:2401-2406 (1993); Mulligan et al., J. Immunol., 150:2407-2417 (1993)], spontaneous colitis in the monkey [Poldolsky et al., J. Clin. Invest., 92:372-380 (1993)] and asthma in  
30 sheep [Lobb, WO 92/13798 published July 22, 1993].

Thus, a preliminary conclusion from in vivo results is that VLA-4 contributes to inflammatory responses that emulate chronic conditions in humans. In an in vivo model of murine contact hypersensitivity, the  
35 CS-1 peptide partially inhibited recruitment of T

- 8 -

lymphocytes to skin inflammatory sites [Ferguson et al.,  
Proc. Natl. Acad. Sci. USA, 88:8072-8076 (1991)].

Because the Arg-Gly-Asp peptide from the cell adhesion  
domain of FN was also inhibitory in this animal model,  
5 the authors concluded that emigration of immune T cells  
to sites of antigenic challenge in the tissue could be  
facilitated by the interaction of leukocyte integrins  
with ECM proteins such as FN [Ferguson et al., Proc.  
Natl. Acad. Sci. USA, 88:8072-8076 (1991)].

10 In a more recent study, Elices and coworkers  
[Elices et al., Clin. Exp. Rheumatol., 11:S77-80 (1993)]  
were unable to reproduce inhibition of contact  
hypersensitivity with the native CS-1 peptide. Instead,  
they found that the CS-1 peptide was rapidly cleared  
15 from blood circulation by proteolytic degradation.


The role of VLA-4 and the CS-1 peptide in  
various chronic and acute immunoinflammatory disease  
states having been established, it would be of  
importance if compounds could be found that inhibit the  
20 VLA-4-lymphocyte interaction and were other than anti-  
VLA-4 antibodies that can themselves induce an immune  
response on repeated administration or the CS-1 peptide  
that is large and costly to make, and also is subject to  
rapid degradation. The disclosure that follows  
25 describes such small molecules that are more potent than  
.is CS-1 itself.

#### Brief Summary of the Invention

30 The present invention contemplates CS-1  
peptidomimetic inhibitor peptides, their compositions  
and methods (processes) for using those inhibitor  
peptides.

- 9 -

A contemplated peptide corresponds in sequence to formula A



5

wherein B is an  $\alpha$ -hydrophobic amino acid residue whose side chain has a length of about 2 to about 5 methylene groups that is preferably leucine;

10 X is a group amide-linked to the nitrogen atom of the B  $\alpha$ -amine. The X group has a ring structure bonded to the carbonyl carbon of the amide-linkage by a spacer having a length of zero to about two methylene groups. The length of X, including the spacer and carbonyl carbon, is about that of a 3-quinolinecarbonyl group or smaller. The ring structure is a 5- and 15 6-membered ring or a fused 6,6- or 6,5-membered ring;


Z is selected from the group consisting of:

(a) Xaa-NCy<sup>1</sup> where Xaa is Val, Ile, Leu or an aromatic amino acid residue; i.e., a residue having a 20 side chain that contains one or two fused aromatic rings, and NCy<sup>1</sup> is a cyclic ring-containing group having a ring nitrogen atom that forms an amide bond with the  $\alpha$ -carboxyl group of Xaa, and whose cyclic ring contains 5- or 6-atoms including said ring nitrogen atom; and

25 (b) NCy<sup>2</sup> where the depicted nitrogen is an amine substituent of a cyclic group whose depicted nitrogen atom forms an amide bond with the  $\alpha$ -carboxyl group of the Asp, and which amine substituent is bonded to a 6- or 7-membered ring or to a fused 6,6- or 30 6,7-membered lactam ring system in which the ring bearing the amine substituent is saturated and contains the amine substituent  $\alpha$  to the carbonyl group of the lactam.

- 10 -

A preferred contemplated peptide corresponds in sequence to formula I



5

wherein X is a group amide-linked to the nitrogen atom of the Leu  $\alpha$ -amine. The X group has a ring structure bonded to the carbonyl carbon of the amide-linkage by a spacer having a length of zero to about two methylene groups. The length of X, including the spacer and carbonyl carbon, is about that of a 3-quinolinecarbonyl group or smaller. The ring structure is a 5- and 6-membered ring or a fused 6,6- or 6,5-membered ring;

15

Z is selected from the group consisting of:

(a) Xaa-NCy<sup>1</sup> where Xaa is Val, Ile, Leu or an aromatic amino acid residue; i.e., a residue having a side chain that contains one or two fused aromatic rings, and NCy<sup>1</sup> is a cyclic ring-containing group having a ring nitrogen atom that forms an amide bond with the  $\alpha$ -carboxyl group of Xaa, and whose cyclic ring contains 5- or 6-atoms including said ring nitrogen atom; and

20

(b) NCy<sup>2</sup> where the depicted nitrogen is an amine substituent of a cyclic group whose depicted nitrogen atom forms an amide bond with the  $\alpha$ -carboxyl group of the Asp, and which amine substituent is bonded to a 6- or 7-membered ring or to a fused 6,6- or 6,7-membered lactam ring system in which the ring bearing the amine substituent is saturated and contains the amine substituent  $\alpha$  to the carbonyl group of the lactam.

25

30


A peptide of formula I, and formulas II and III hereinafter, is water-soluble and inhibits the binding of Jurkat cells (ATCC TIB 152) to a solid phase-bound peptide of SEQ ID NO:1 in an in vitro assay in an

35

- 11 -

aqueous buffer at a pH value of 7.2-7.4 to an extent that is about 10- to about 3000-fold better than the inhibition in the binding exhibited by a peptide of SEQ ID NO:3.

5 In a more preferred embodiment, the X group has the formula Ar-Y-C(O)- and the peptide corresponds in sequence to formula II, below,



10

wherein Ar is a pyrazolyl, phenyl, pyridyl, or 3-quinolinyl group;

Y is a spacer that is absent, -CH<sub>2</sub>-, -CH(NH)-, -O- or -NH-;

15

or Ar-Y-C(O)- together with the nitrogen atom of said Leu forms a phthalimido, a 1,2,3,4-tetrahydroquinazoline-2,4-dione-3-yl or a 5-phenyldantoin-3-yl group, and

20

Ar-Y-C(O)- has a length of about 3-quinolinecarbonyl or less.


Xaa is an aromatic amino acid residue; i.e., an amino acid residue having an aromatic side chain.

25

NCy<sup>1</sup> is an amine-containing 5- or 6-membered cyclic ring group whose depicted nitrogen atom is within the ring and forms an amide bond with α-carboxyl of Xaa, as discussed before.

Ar-Y-C(O)- is most preferably phenylacetyl, so that in most preferred practice, a contemplated peptide corresponds in sequence to formula III, below,

30



35

wherein NCy<sup>3</sup> is a most preferred NCy<sup>1</sup> group that is selected from the group consisting of morpholinamido, thiomorpholinamido,

- 12 -

4-(thiadioxo)piperidinamido, D-2-(carboxamide)-  
pyrrolidinamido, piperidinamido and a substituted  
piperidinamido whose substituent is selected from the  
group consisting of 4-hydroxy, 4-carbamyl and 4-carboxy,  
5 piperazinamido, substituted piperazinamido whose  
substituent is selected from the group consisting of  
4-N-carboxymethyl, 4-N-carboxamidomethyl, 4-N-(5-  
hydroxyethylenoxyethylene) and pyrrolidinamido groups.

A composition containing an above inhibitor  
10 peptide is also contemplated. Such a composition  
contains the peptide dissolved or dispersed in a  
pharmaceutically acceptable diluent that is preferably  
aqueous. The peptide is present in the composition in  
an inflammation-reducing amount. That amount is also  
15 sometimes referred to herein as a CS-1/VLA-4 binding  
inhibiting amount. An above-discussed more preferred or  
most preferred peptide is utilized in a more preferred  
or most preferred composition.

A process for treating fibronectin CS-1/VLA-4-  
20 mediated inflammation is also contemplated. That  
process comprises administering to a mammal having that  
inflammation or otherwise in need of such a treatment as  
for prophylactic purposes, an inflammation-reducing  
amount of a before-described inhibitor peptide. Use of  
25 a more preferred or most preferred inhibitor peptide is  
more or most preferred in this process.

All peptide formulas or sequences shown herein  
are written from left to right and in the direction from  
amino-terminus to carboxy-terminus. The abbreviations  
30 used herein for derivatives and residues of the twenty  
natural amino acids are reproduced in the following  
Table of Correspondence:

- 13 -

TABLE OF CORRESPONDENCE

	<u>Abbreviation</u>		<u>Amino Acid</u>
	<u>1- Letter</u>	<u>3-Letter</u>	
	Y	Tyr	tyrosine
5	G	Gly	glycine
	F	Phe	phenylalanine
	M	Met	methionine
	A	Ala	alanine
	S	Ser	serine
10	I	Ile	isoleucine
	L	Leu	leucine
	T	Thr	threonine
	V	Val	valine
	P	Pro	proline
15	K	Lys	lysine
	H	His	histidine
	Q	Gln	glutamine
	E	Glu	glutamic acid
	W	Trp	tryptophan
20	R	Arg	arginine
	D	Asp	aspartic acid
	N	Asn	asparagine
	C	Cys	cysteine
25	X	Xaa	another residue, or one of several residues

The present invention has several benefits and  
30 advantages.

One salient benefit is that an inhibitor  
peptide contemplated here is more potent in inhibiting  
the VLA-4/CS-1 binding interaction than CS-1 itself.

Another benefit of the invention is that  
35 exemplary inhibitor peptides have been shown to be

- 14 -

effective in reducing various exemplary immunoinflammation disease states in host mammals.

An advantage of the invention is that no adverse toxicity has been observed at in vivo concentrations of up to 600 mg/kg/day in a host laboratory mammal.

Another advantage of the invention is that a contemplated inhibitor peptide is a relatively small molecule that is easily prepared in high yield and purity.

Still further benefits and advantages of the invention will become apparent to the skilled worker from the disclosure that follows.

#### Brief Description of the Drawings

In the drawings forming a portion of this disclosure:

Fig. 1 is a graph illustrating the in vitro binding inhibition of VLA-4-bearing Jurkat cells to the solid phase-bound CS-1 peptide (SEQ ID NO:1) by that peptide itself and shorter peptides having portions of the CS-1 peptide sequence. Data are shown as percentages relative to the indicated "Standard" (SEQ ID NO:3). Data for peptides with deletions at the "N-terminus" of peptide B12 (SEQ ID NO:2) are shown to the left of the Standard, and data for peptides with deletions at the "C-terminus" of peptide B12 are shown to the right of the standard. Peptide sequences are in single letter code.

Fig. 2 is a graph with data obtained and expressed similarly to those of Fig. 1. Here, binding inhibition by further, still shorter deletion peptides, is illustrated.

Fig. 3 is another graph of binding data obtained as discussed in Fig. 1. This graph utilizes an



- 15 -

ordinate that is on a log scale. These data are arranged into five groups and are again shown relative to the indicated Standard peptide (SEQ ID NO:3), with D-proline being shown as "p". The data of the right-hand-most two groups illustrate the effects of three different X groups on a single indicated peptide and of three Z groups in which X is phenylacetyl ( $\phi$ Ac) and Z is as shown, respectively.

Fig. 4 shown in two panels as Fig. 4A and Fig. 4B illustrates the effects of a contemplated peptide in treating asthma in the rabbit. Fig. 4A shows the percent change in dynamic compliance ( $C_{dyn}$ ) over a six-hour time period immediately following the onset of induced asthma attacks. Data for rabbits treated by a nebulized composition containing the inhibitor peptide N-phenylacetyl-Leu-Asp-Phe-morpholinamide are shown as open circles, whereas data for untreated rabbits are shown with darkened circles; both circles including error bars. The ordinate is in units of percent change from the initial dynamic compliance value, whereas the abscissa is in units of hours after challenge.

Fig. 4B shows results obtained for the percent change in lung resistance ( $L_R$ ) from the same study, with data being presented as in Fig. 4A. The ordinate is in units of percent change from the original lung resistance value, whereas the abscissa is in hours.

Fig. 5 is a graph showing the results of a study of the effect of the inhibitor peptide N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub> on delayed type hypersensitivity measured in ears of 14 mice. After immunization, one group of seven mice was treated with only a saline solution provided by an implanted pump over a 24-hour time period and challenged. The other immunized group of seven mice was similarly challenged, but each was treated with an aqueous pharmaceutical

- 16 -

composition containing the above inhibitor peptide for the same 24-hour time period, also supplied by implanted pumps. The ordinate is in units of inches of swelling diameter at the challenge site.

5                   Fig. 6 is a graph showing averaged clinical scores for six mice each in two evaluations of treatments of experimental autoimmune encephalomyelitis (EAE). Darkened circles are for treatments using the inhibitor peptide N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub>,  
10                   whereas points shown as darkened squares are for treatments using the scrambled sequence peptide of Example 6. The ordinant shows the averaged score for the six mice in each study, whereas the abscissa is in days after initiation of EAE.

15                   Fig. 7 is a graph showing the percentage of change in lung resistance, SR<sub>L</sub>, from base line for the asthmatic sheep model depicted as described in Fib. 4B. The nebulized composition here contained N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub> in open circles and the scrambled  
20                   sequence of that peptide (Example 6) in darkened circles, including error bars where appropriate.

#### Detailed Description of the Invention

25                   The present invention contemplates an inhibitor peptide, a composition containing such a peptide, and a process of using such a peptide. A contemplated peptide inhibits binding between the CS-1 peptide of fibronectin and the inflammatory cell VLA-4 surface receptor, and is therefore sometimes referred to  
30                   herein as an inhibitor peptide.

##### A. Peptides

35                   A contemplated peptide can be viewed as a mimic of the whole fibronectin molecule, or at least the 25-residue (25-mer) CS-1 portion of fibronectin (SEQ ID NO:1) that binds to the VLA-4 receptor. As will be seen

- 17 -

from the discussion that follows, a contemplated peptide binds to the VLA-4 receptor even more tightly than does the CS-1 25-mer peptide present in fibronectin.

5 Broadly, a contemplated inhibitor peptide can be defined as having a structure corresponding to formula A, below



10 wherein

B is an  $\alpha$ -hydrophobic amino acid residue whose side chain has a length of about 2 to about 5 methylene groups;

15 X is a group amide-linked to the nitrogen atom of the B  $\alpha$ -amine. The X group has a ring structure bonded to the carbonyl carbon of the amide-linkage by a spacer having a length of zero to about two methylene groups. The length of X, including the spacer and carbonyl carbon, is about that of a 3-quinolinecarbonyl group or smaller. The ring structure is a 5- and 20 6-membered ring or a fused 6,6- or 6,5-membered ring;

Z is selected from the group consisting of:

(a) Xaa-NCy<sup>1</sup> where Xaa is Val, Ile, Leu or an aromatic amino acid residue; i.e., a residue having a 25 side chain that contains one or two fused aromatic rings, and NCy<sup>1</sup> is a cyclic ring-containing group having a ring nitrogen atom that forms an amide bond with the  $\alpha$ -carboxyl group of Xaa, and whose cyclic ring contains 5- or 6-atoms including said ring nitrogen atom; and

30 (b) NCy<sup>2</sup> where the depicted nitrogen is an amine substituent of a cyclic group whose depicted nitrogen atom forms an amide bond with the  $\alpha$ -carboxyl group of the Asp, and which amine substituent is bonded to a 6- or 7-membered ring or to a fused 6,6- or 35 6,7-membered lactam ring system in which the ring

- 18 -

bearing the amine substituent is saturated and contains the amine substituent  $\alpha$  to the carbonyl group of the lactam.

5 A peptide of formula A is water-soluble and inhibits the binding of Jurkat cells (ATCC TIB 152) to a solid phase-bound peptide of SEQ ID NO:1 in an in vitro assay in an aqueous buffer at a pH value of 7.2-7.4 to an extent that is equal to or up to about 3000-fold greater than the binding inhibition exhibited by a  
10 peptide of SEQ ID NO:3.


The sequence of a peptide of formula A appears to require the Asp residue present in the CS-1 (SEQ ID NO:1) and B12 (SEQ ID NO:3) fibronectin peptides. Aside from that Asp, both of whose size and charge appear to  
15 be required for binding as Glu and other residues barely inhibit binding, size and relative hydrophobicity appear to be most important in the selection of the B residue. The requirements of the other residues are discussed hereinafter.

20 The length of a B residue side chain is about 2 to about 5 methylenes. Such lengths can be readily determined from standard bond lengths, Van der Waals radii, by use of commercially available atomic models or computer programs. Using such lengths, it is seen that  
25 both cyclohexylalanine and phenylalanine have a side chain length of about 5 methylenes; i.e., one methylene plus four ring carbons in the side chain. A methionine side chain, having a methyl, thio and two methylenes, is shorter still.

30 Exemplary B residues as amino acids are selected from the group consisting of leucine (Leu), cyclohexylalanine, norleucine (Nle), Methionine (Met), homoserine, threonine (Thr), phenylalanine (Phe), valine (Val), norvaline (Nva), and isoleucine (Ile). B is most  
35 preferably Leu.

- 19 -

Preferably, a contemplated inhibitor peptide has a structure corresponding to formula I, below



5

wherein

X is a group amide-linked to the nitrogen atom of Leu, the group having a ring structure bonded to the carbonyl carbon of the amide-linkage by a spacer having  
10 a length of zero to about two methylene groups. The length of X, including the spacer and carbonyl carbon, is about that of a 3-quinolinecarbonyl group or smaller. The ring structure is a 5- and 6-membered ring or a fused 6,6- or 6,5-membered ring; and

15

Z is selected from the group consisting of

(a) Xaa-NCy<sup>1</sup> where Xaa is Val, Ile, Leu or an amino acid residue having a side chain that contains one or two fused aromatic rings and NCy<sup>1</sup> is a cyclic ring-containing group having a ring nitrogen atom that forms  
20 an amide bond with the  $\alpha$ -carboxyl group of Xaa, and whose cyclic ring contains 5- or 6-atoms including said ring nitrogen atom; and

25

(b) NCy<sup>2</sup>, where the depicted nitrogen is an amine substituent of a cyclic group whose depicted nitrogen atom forms an amide bond with the  $\alpha$ -carboxyl group of the Asp residue, and which amine substituent is bonded to a 6- or 7-membered ring or to a fused 6,6- or 6,7-membered lactam ring system in which the ring bearing the amine substituent is saturated and contains  
30 the amine substituent  $\alpha$  to the carbonyl group of the lactam.

35

A peptide of formula I, as well as formulas II and III below, is water-soluble and inhibits the binding of Jurkat cells to a solid phase-bound peptide of SEQ ID NO:1 in an in vitro assay in an aqueous buffer at a pH

- 20 -

value of 7.2-7.4 to an extent that is about 10- to about 3000-fold better than the inhibition in said binding exhibited by a peptide of SEQ ID NO:3.

Examining formula I, it is seen that at least  
5 Leu and Asp of the CS-1 (SEQ ID NO:1) and B12 (SEQ ID NO:2) fibronectin peptides are present. Aside from that two residue sequence, the sequence/structure of a contemplated inhibitor peptide and the CS-1 or B12 portions are quite different.

10 Thus, whereas there is an iso-leucine (Ile;I) peptide- (amide-) bonded to the  $\alpha$ -amine group of the Leu residue in CS-1 and the native protein, a cyclic ring structure-containing group or moiety, X, is amide-bonded to the nitrogen atom of the B or Leu residue  $\alpha$ -amino  
15 group (formulas A or I, respectively) via a carboxyl contributed by the cyclic ring structure-containing group. That amide bond can be present as part of a carboxamide- [-C(O)NH-], urethane- [-O-C(O)NH-] or urea- [-NH-C(O)NH-] containing spacer group that links the  
20 cyclic ring structure-containing group to the Leu residue.

The cyclic ring structure broadly can be any 5- or 6-membered ring that is saturated or contains ethylenic unsaturation. The ring structure can contain  
25 one or more atoms other than carbon such as nitrogen, oxygen or sulfur. The ring structure can also be a fused ring system where two 6-membered rings are fused (6,6-) or where a 6-membered ring is fused to a 5-membered ring (6,5-membered). The ring of the cyclic  
30 ring structure is preferably aromatic.

Exemplary ring structures include tetrahydrofuryl, tetrahydropyryl, cyclopentyl, cyclohexyl, phenyl, pyridyl, pyrimidinyl, pyrazinyl, pyrazolyl, pyrrolidyl, furanyl, piperidinyl, naphthyl,  
35 quinolinyl, decalinyl, quinazolinyl, imidazolyl,

- 21 -

thiophenyl, and the like. Of the cyclic ring structures, phenyl and pyridyl are particularly preferred.

5 A cyclic ring structure can be bonded directly to the carbonyl group [-C(O)-] of the amide bond to the B or Leu residue. That ring can also be spaced away from the carbonyl group by up to about the length of two methylene (-CH<sub>2</sub>-) groups or an ethylene group (-CH<sub>2</sub>-CH<sub>2</sub>-).

10 The Van der Waals radius a methylene group (about 2.0 Å) is slightly longer than that of an oxy group (-O-; about 1.40 Å) or an imino group (-NH-; about 1.50 Å). There is sufficient similarity between the sizes of methylene, oxy and imino so that a spacer group  
15 containing a -CH<sub>2</sub>-O-, -CH<sub>2</sub>-NH-, -NH-NH-, or -O-NH- are of similar lengths and are within the length of an ethylene group, -CH<sub>2</sub>-CH<sub>2</sub>-. A similar result obtains if bond lengths (distances) are used. Contemplated spacers include -HC(CH<sub>3</sub>)-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-, -NH-O-, -HN-NH-, -CH<sub>2</sub>-O-  
20 and -CH<sub>2</sub>-NH-, and are preferably free of unsaturation.

Using a phenyl ring as an exemplary aromatic ring structure, it is seen that the contemplated X groups include 3-methyl-3-phenylpropionyl, 3-phenylpropionyl, phenylhydroxaminocarbonyl [Ph-NH-O-C(O)-],  
25 phenylhydrazidecarbonyl [Ph-NH-NH-C(O)-], benzyloxycarbonyl [Ph-CH<sub>2</sub>-O-C(O)-], phenoxyacetyl [Ph-O-CH<sub>2</sub>-C(O)-], benzylaminocarbonyl [Ph-CH<sub>2</sub>-NH-C(O)-], and anilinoacetyl [Ph-NH-CH<sub>2</sub>-C(O)-], where "Ph" is a phenyl group.

30 Thus, it is contemplated that a before-described ring structure be bonded to the carbonyl carbon of the B- or Leu-linked amide group by a spacer having a length of zero methylene groups (a direct bond), one or two methylene groups. Put differently,

- 22 -

the spacer has the length of about an ethylene group or less.

A phenylacetyl, phenoxycarbonyl or anilinocarbonyl group bonded to the nitrogen of the B or Leu  $\alpha$ -amino group contains a spacer having the length of about one methylene group. Phenyl (benzoyl), 1- or 2-naphthyl (1- or 2-naphthalenecarbonyl), 2-, 3- or 4-pyridyl (2-, 3- or 4-pyridinecarbonyl), 2- or 3-thiophenyl (2- or 3-thienyl; 2- or 3-thiophencarbonyl) and 2- or 3-furanyl (2- or 3-furancarbonyl) ring structures are bonded directly to the amide carbonyl carbon and therefore define an X group that utilizes a spacer having a length of zero methylene groups. A spacer having a length of about two methylene groups is provided by an X group that is carbobenzyloxy [ $\text{Ph-CH}_2\text{-O-C(O)-}$ ], carbobenzylamino [ $\text{Ph-CH}_2\text{-NH-C(O)-}$ ], carbophenoxymethylene [ $\text{Ph-O-CH}_2\text{-C(O)-}$ ] and the like groups.

A contemplated 5- or 6-membered ring structure can also be substituted with a  $\text{C}_1\text{-C}_2$  alkyl or hydroxyl group. Exemplary substituted ring structures using a phenyl ring as illustrative thus include 2-, 3- or 4-ethylphenyl, 2,6-, 3,4- or 2,3-dimethylphenyl, 2-, 3- or 4-hydroxyphenyl, 2,6-, 2,4-, 3,4- and 3,5-dihydroxyphenyl, and the like.

The ring structure of the X substituent is thought to act in a contemplated inhibitor in some way to fit the inhibitor peptide into the binding pocket of the VLA-4 receptor to position the B or Leu and Asp groups into a proper configuration. Because of that presumed role in fitting the peptide into its receptor, there are some size constraints upon the ring structure-containing and spacer portions of X, in addition to those noted before as to the spacer group length. Thus, from the carbonyl-containing carbon of the amide bond to



- 23 -

B or Leu, through the end of ring structure or its substituent furthest from the carbonyl group, the total length of the spacer plus ring structure-containing portion of X is about the size of a 3-quinolinecarbonyl group or smaller.

Inasmuch as a 3-quinolinecarbonyl group is the longest contemplated ring structure-containing X substituent, a 3-quinolinecarbonyl group is free from the above-discussed substituents that add to its length.

The length of a given X substituent can be readily determined, as discussed before. For example, one can use space-filling models to build exemplary cyclic ring structure-containing X groups and then compare the relative sizes of the prepared models. One can also use published bond lengths and bond angles to prepare a two-dimensional depiction of the sizes. Computer graphics programs are also well-known and available that can be used to prepare exemplary model X groups for length comparison to 3-quinolinoyl.

The X substituent, including the spacer, cyclic ring structure, the carbonyl group and the  $\alpha$ -amino nitrogen atom of B or Leu can also together form an aromatic ring-substituted cyclic imido group. Exemplary of such cyclic imido groups are phthalimido, which is preferred, each of 2,3- and 3,4-pyridinedicarboximido, homophthalimido and 1,2,3,4-tetrahydroquinazoline-2,4-dione-3-yl groups in which the aromatic ring and cyclic imido group are fused together.

In another exemplary compound, the B or leucine nitrogen atom is an imido nitrogen atom within the ring of a 5-phenylhydantoin-3-yl group so that the aromatic phenyl ring is a substituent of a cyclic spacer and is spaced about one methylene away from the carbonyl group linked to the Leu residue. A similarly structured imido nitrogen-containing X group is present in a

- 24 -

2-phenylsuccinimido group formed on the B or Leu nitrogen atom.

5 The cyclic imido- and hydantoin-containing portions of the above-discussed X groups can thus be viewed as specialized spacer groups that limit the conformational degrees of freedom of the ring structures. Thus, for example, whereas the carbonyl, methylene and phenyl portions of a phenylacetyl group are each free to assume one or more of several  
10 conformations, a phthalimido X group can only spin about the axis of the leucine nitrogen-methine bond.

It is noted that although the X substituent must contain a cyclic ring structure that can be substituted as discussed before, that X substituent can  
15 also include a further substituent on other than the ring structure. When a further substituent is present, X preferably is an amino acid residue having a cyclic ring side chain that therefore includes a primary or secondary amine. Here, X is preferably a prolyl, phenylalanyl, tyrosinyl or phenylglycyl residue, the  
20 nitrogen atom of whose  $\alpha$ -amino group is bonded to the further substituent.

That further substituent can be one amino acid residue through the remainder of the CS-1 polypeptide  
25 sequence toward the N-terminus thereof, with the sequence of that polypeptide beginning at the isoleucine of position 19 from the N-terminus of SEQ ID NO:1. A single residue or 18 separate amino acid residue substituent sequences are thereby defined.

30 Another exemplary further substituent linked via an amine group of X is biotin. In a particular example, biotin amide-bonded to  $\epsilon$ -aminocaproic acid was amide-bonded to the  $\alpha$ -amine of a phenylalanine (Phe) as an X group. The resulting compound contained the biotin

- 25 -

fused ring amide-linked to the Phe X group via a chain of twelve atoms.

It should also be understood that an X group amino acid residue having a cyclic ring side chain can also be free of substituent groups. The nitrogen atom of the  $\alpha$ -amine of such a residue can also be acylated as with a  $C_1$ - $C_6$  acyl group such as formyl, acetyl, iso-butyryl, or hexanoyl group. A  $C_1$ - $C_6$  acyl group bonded to the nitrogen of an  $\alpha$ -amine group forms an amide bond at that nitrogen atom and provides no ionic charge to the peptide at a pH value of 7.2-7.4 as compared to the positive charge provided by an unsubstituted free  $\alpha$ -amine.

The Z group of a before-discussed formula can be one of two types of groups. The Z group in one embodiment (A) is a hydrophobic amino acid residue Xaa peptide-bonded to the Asp carboxyl and linked to a cyclic ring-containing group  $NCy^1$  that has a ring nitrogen atom (the N of  $NCy^1$ ) that forms an amide bond with the  $\alpha$ -carboxyl group of Xaa. The cyclic ring of  $NCy^1$  contains 5- or 6-atoms, including the depicted nitrogen atom (N of  $NCy^1$ ). Contemplated hydrophobic amino acid residues are those having aliphatic side chains such as valine, leucine and isoleucine. Xaa more preferably contains a hydrophobic aromatic amino acid residue; i.e., Xaa is an amino acid residue having an aromatic side chain that contains one or two fused aromatic rings. Exemplary of such aromatic amino acids are phenylalanine, tyrosine and tryptophan that are naturally occurring (genetically encoded) as well as phenylglycine, homophenylalanine, p-nitrophenylalanine, thiophenylglycine (thienylglycine), and the like.

Exemplary  $NCy^1$  groups include morpholinyl, thiomorpholinyl, thiomorpholinylsulfone [4-(thiadioxo)piperidinyl], piperidinyl, piperazinyl,

- 26 -

pyrrolidinyl, pyrazolyl, pyrazolinyl, oxazolidinyl and the like as their respective amides. A NCy<sup>1</sup> cyclic ring can also be substituted with one or two substituent groups selected from the group consisting of carboxyl, carboxamide, C<sub>1</sub>-C<sub>4</sub> alkylenecarboxyl C<sub>1</sub>-C<sub>4</sub> alkylenecarboxamide, hydroxyl, hydroxymethyl, (CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>H where n is one, two or three and C<sub>1</sub>-C<sub>4</sub> alkyl. Carboxyl substitution at the 2-position of a pyrrolidine provides the amino acid proline, whose D- and L-forms are both contemplated herein. D-Prolyl (sometimes shown in bold face lower case single letter amino acid code as "p" or as D-Pro) is particularly preferred as its amido derivative (D-Pro-NH<sub>2</sub>) as are morpholinyl, piperidyl, piperazinyl and 4-hydroxypiperidyl.

Exemplary C<sub>1</sub>-C<sub>4</sub> alkyl groups include methyl, ethyl, iso-propyl, n-butyl and t-butyl. A C<sub>1</sub>-C<sub>4</sub> alkyl group can also form a quaternary ammonium group with a second nitrogen atom of NCy<sup>1</sup> such as piperazine. Where the C<sub>1</sub>-C<sub>4</sub> alkyl group is a methyl group, and iodide is the anion, and exemplary NCy<sup>1</sup> group is a 4,4-N,N-dimethylpiperaziniumyl iodide.

Exemplary C<sub>1</sub>-C<sub>4</sub> alkylenecarboxyl and C<sub>1</sub>-C<sub>4</sub> alkylenecarboxamide groups include methylenecarboxyl (-CH<sub>2</sub>CO<sub>2</sub>H; carboxymethyl) and methylenecarboxamido (-CH<sub>2</sub>CONH<sub>2</sub>; carboxamidomethyl), ethylenecarboxyl (-CH<sub>2</sub>CH<sub>2</sub>CO<sub>2</sub>H; carboxyethyl) and ethylenecarboxamido (-CH<sub>2</sub>CH<sub>2</sub>CONH<sub>2</sub>; carboxamidoethyl) as well as butylenecarboxyl (-C<sub>4</sub>H<sub>9</sub>CO<sub>2</sub>H; carboxybutyl) and butylenecarboxamido (-C<sub>4</sub>H<sub>9</sub>CONH<sub>2</sub>; carboxyamidobutyl).

Exemplary groups (CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>H where n is one, two or three include 2-hydroxyethyl (n=1), 5-hydroxyethylenoxyethylene (ethyleneoxyethanol; 5-hydroxy-3-oxapentyl; n=2) and 8-hydroxy-3,5-dioxa-octyl (n=3).

When NCy<sup>1</sup> includes a piperazinyl group, the second (4-position) nitrogen atom cannot only be

- 27 -

quaternized by alkylation, but also amidified. Exemplary acyl portions of the piperazinyl-4-N-amides include C<sub>1</sub>-C<sub>6</sub> acyl groups such as formyl, acetyl, propionyl, isobutanoyl, hexanoyl and benzoyl, but also sulfonamides such as phenylsulfonamido, toluenesulfonamide (tosyl), methanesulfonamide (mesyl) and trifluoromethylsulfonamido (trifyl).

Thus, in those embodiments where Z is Xaa-NCy<sup>1</sup>, Xaa is a specified amino acid residue whose amine group forms an amide (peptide) bond with the  $\alpha$ -carboxyl of the depicted Asp residue, and whose carboxyl group forms an amide bond with a nitrogen atom present within the 5- or 6-membered ring of NCy<sup>1</sup>.

In another embodiment (B) Z is NCy<sup>2</sup> where the depicted nitrogen atom (N of NCy<sup>2</sup>) is an amine substituent of a cyclic group (Cy<sup>2</sup>) whose substituent nitrogen atom forms an amide bond with the  $\alpha$ -carboxyl of the depicted Asp residue. That amine substituent is bonded to a cyclic group that is (i) a 6- or 7-membered ring or (ii) a fused 6,6- or 6,7-membered lactam ring system in which the ring bearing the amine substituent is saturated (free of ethylenic unsaturation) and contains the amine substituent  $\alpha$  to the carboxyl group of the lactam.

Here, the nitrogen atom that links the ring system to the remainder of the peptide is a substituent of a cyclic ring structure rather than being a ring atom as in NCy<sup>1</sup>. In addition, the rings of which that nitrogen can be a substituent are of two types, 6- or 7-membered rings or 6,6- or 6,7-membered fused ring systems, one of which rings is a lactam. In either situation, there is no Xaa amino acid residue in this embodiment.

Exemplary amine substituent-containing 6- and 7-membered ring NCy<sup>2</sup> groups of this type include

- 28 -

benzylamine, phenethylamine, 2-(N-morpholinyl)-ethylamine, N-[1-(carboxamidomethyl)-caprolactam-3-yl]amine, N-(caprolactam-3-yl)amine, and N-(valerolactam-3-yl)amine groups that form the corresponding amides with the  $\alpha$ -carboxyl of Asp. Exemplary amino-substituted 6,6- and 6,7-fused ring lactam-containing NCy<sup>2</sup> groups include N-[1-(2-N-morpholinylethyl)-2-oxo-tetrahydroquinolin-3-yl]amine, N-(2-oxo-tetrahydroquinolin-3-yl)amine and the 6,7-fused ring tricyclic compound shown at footnote 7 of Table 1 groups that form corresponding amides with the  $\alpha$ -carboxyl of Asp.

It has generally been found that once (i) the X group of a formula discussed herein is occupied by an aromatic ring-containing moiety spaced adjacent to or within about one methylene group's distance from the carbonyl, (ii) Z is an aromatic amino acid, and (iii) NCy<sup>1</sup> is L- or D-proline amide or a 5- or 6-membered nitrogen-containing ring as discussed before, substantially any other substituent can be present linked to either peptide terminus without abolishing the inhibitory activity of a contemplated peptide, so long as the resulting peptide is water-soluble.

Thus, for example, the peptide of SEQ ID NO:5 having an N-terminal phenylacetyl group linked to the sequence Leu-Asp-Phe-Pro can further include a substituted tetraethylenediamine group amide-bonded to the Pro residue in which four phenylacetyl-Leu-Asp-Phe-Pro groups were amide-bonded to the tetraethylenediamine nitrogens and still exhibit VLA-4 binding inhibition that was about 210 times better than the standard 10-mer peptide of SEQ ID NO:3.

Similarly, the peptide PheLeuAspPhe-D-Pro-NH<sub>2</sub> contained a europium-containing chelate at its N-terminus bonded to the nitrogen atom of the N-terminal

- 29 -

Phe. That peptide exhibited a binding inhibition about 120 times better than that of the peptide of SEQ ID NO:3.

5 The peptide of SEQ ID NO:12, phenylacetyl-Leu-  
Asp-Phe-Pro-NH(CH<sub>2</sub>)<sub>5</sub>C(O)NHC<sub>18</sub>H<sub>37</sub>, would be predicted to be  
a good inhibitor. However, that peptide is not water-  
soluble and forms a turbid dispersion rather than a  
solution. That peptide exhibits a binding inhibition  
similar to that exhibited by the standard 10-mer peptide  
10 of SEQ ID NO:3.

Binding phenomena involving structure-function  
relations such as those contemplated herein are not  
completely understood, although a skilled worker can  
usually make accurate predictions as to desired potency  
15 of a predicted peptide once sufficient data are in hand  
as is discussed hereinafter. Indeed, substantially all  
of the compounds defined by a before discussed formula  
inhibit binding between the CS-1 peptide and the VLA-4  
receptor. Nonetheless, because the substituents bonded  
20 directly to Leu and Asp of formula I can vary widely and  
affect binding inhibition to VLA-4, a preferred  
contemplated inhibitor peptide is defined not only by  
its structure, but also by its potency as an inhibitor  
in a standard in vitro assay as compared to a standard,  
25 known, 10-mer peptide inhibitor, to include only those  
inhibitor peptides that inhibit binding by about one  
order of magnitude or more better than the known 10-mer  
inhibitor.

A contemplated inhibitor peptide thus inhibits  
30 the binding of inflammatory cells that contain the VLA-4  
receptor [Jurkat cells (American Type Culture  
Collection, Rockville, MD 20852, ATCC TIB 152)] to the  
solid phase-bound CS-1 peptide (SEQ ID NO:1) in an  
aqueous buffer at pH 7.2-7.4 to an extent that is about  
35 10-fold to about 1000-fold and more preferably 3000-fold


- 30 -

better than that binding exhibited by the art standard 10-mer peptide of SEQ ID NO:3 [GPEILDVPST in single letter code). More preferably, that binding is inhibited by about 50- to about 3000-fold, and most preferably by about 100- to about 3000-fold.

Binding inhibition is measured here as a concentration of peptide that inhibits one-half the binding between a standard number of Jurkat cells and a standard amount of CS-1 peptide bound to the surface of a microtiter plate well. Those concentrations are conveniently expressed as  $IC_{50}$  values, smaller numbers indicating a lower concentration required to inhibit 50 percent binding and therefore greater potency. Further specifics of this assay are provided hereinafter.

To recapitulate, a peptide of formulas A or I inhibits binding between the CS-1 peptide region of fibronectin and the VLA-4 receptor. However, those inhibitors that are at least ten-times better inhibitors than the peptide of SEQ ID NO:3 are preferred here.

Still more preferred is a peptide of formula II, below,



wherein Ar is a pyrazolyl, phenyl, pyridyl (2-, 3- or 4-), or 3-quinolinyl group;

Y is a spacer that is absent,  $-\text{CH}_2-$ ,  $-\text{CH}(\text{NH})-$ ,  $-\text{O}-$  or  $-\text{NH}-$ ;

or Ar-Y-C(O) together with the nitrogen atom of Leu forms a phthalimido, a 1,2,3,4-tetrahydroquinazoline-2,4-dione-3-yl or 5-phenylhydantoin-3-yl group,

Ar-Y-C(O)- has a length of about 3-quinolinecarbonyl or less;



- 31 -

Xaa is an aromatic amino acid residue; i.e., an amino acid residue having an aromatic side chain, such as phenylalanine, tyrosine, tryptophan, homophenylalanine, nitrophenylalanine, thienylglycine and phenylglycine; and

NCy<sup>1</sup> is an amine-containing 5- or 6-membered cyclic ring group whose depicted nitrogen atom, N of NCy<sup>1</sup>, is within the ring and forms an amide bond with the  $\alpha$ -carboxyl of Xaa, as was discussed before.

The water-soluble compounds of formula II thus far examined exhibit binding inhibitions of at least 10-fold better than that exhibited by the peptide of SEQ ID NO:3, and thus no in vitro assay is required for their complete definition, although that assay applies to all of the preferred inhibitor peptides.

Of the above combinations, Ar-Y-C(O), a more preferred X group of formula I, is preferably benzoyl, phenylacetyl, 4-pyridinecarbonyl (isonicotinoyl), 3-pyridinecarbonyl (nicotinoyl), 3-pyridinacetyl, anilino carbonyl, 3-quinolinoyl, pyrazolecarbonyl, tryptophyl and 3,4-dihydroxybenzoyl, with phenylacetyl (benzylcarbonyl) being most preferred, or Ar-Y-C(O) together with the leucine nitrogen atom form a phthalimido group. Xaa is preferably Phe, Tyr or Trp, with Phe being most preferred. NCy<sup>1</sup> is preferably an amide of a morpholinyl, piperidinyl or substituted piperidinyl where the substituent is selected from the group consisting of hydroxyl, carboxyl, carboxamido groups, piperazinyl or 4-substituted piperazinyl in which the 4-substituent is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>1</sub>-C<sub>4</sub> alkylencarboxyl, C<sub>1</sub>-C<sub>4</sub> alkylencarboxamide, (CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>H where n is 1, 2 or 3, thiomorpholinyl, L- or D-prolinyl amide, pyrrolidinyl, 3,4-dihydropyrrolidinyl, 2-(hydroxymethyl)pyrrolidinyl and 4-(thiadioxo)piperidinyl group, with amides of

- 32 -

morpholinyl, D-prolinyl amide, piperidinyl, an above-substituted piperidinyl, piperazinyl, an above-substituted piperazinyl and pyrrolidinyl groups being most preferred.

5                   A most preferred peptide corresponds in sequence to a peptide of formula III, below,



10                   wherein NCy<sup>3</sup> is a group of most preferred NCy<sup>1</sup> groups and is selected from the group consisting of morpholinamido, thiomorpholino, 4-(thiadioxo)piperidinamido, D-2-(carboxamido)pyrrolidinamido, piperazinamido, 15 substituted piperazinamido where the substituent is selected from the group consisting of 4-N-carboxymethyl, 4-N-carboxamidomethyl, 4-N-(5-hydroxyethylenoxyethylene) and 4-N-p-toluenesulfonamido, pyrrolidinamido, piperidinamido and substituted piperidinamido where the 20 substituent is selected from the group consisting of 4-hydroxy, 4-carbamyl, 4-carboxyl groups.

                  Exemplary inhibitor peptides are listed in Table 1 below along with their binding inhibition potencies relative to the standard peptide of SEQ ID 25 NO:3.

TABLE 1  
RELATIVE PEPTIDE INHIBITION OF VLA-4 BINDING  
CS-1

SEQ. ID. NO.	FORMULA'	X	Z'	RELATIVE POTENCY'
		phenylacetyl	4-N(carboxymethyl)piperazinamide'	2579
X L D F Z		phenylacetyl	4-(thiadioxo)piperidinamide'	1060
X L D F Z		phenylacetyl	4-(carbamy)l)piperidinamide''	933
X L D F Z		phenylacetyl	morpholinamide	844
X L D F Z		phenylacetyl	4-(carboxy)l)piperidinamide	828
X L D F Z		pyridine-4-carbonyl	amide	656
X L D F P		phenylacetyl	amide	640
X L D Y P		phenylacetyl	4-hydroxypiperidinamide	603
X L D F Z		phenylacetyl	piperazinamide	506
X L D F Z		phenylacetyl	4-N-(5-hydroxyethyloxyethylene)-piperazinamide''	505
X L D F Z		phenylacetyl	thiomorpholinamide	493
X L D F Z		pyridine-3-acetyl	amide	469
X L D F P		phenylacetyl	amide	405
X L D F P		phenylacetyl	4-N-(p-toluenesulfonamido)-piperidinamide	360
X L D F Z		phenylacetyl	4-N-(carboxamidomethyl)piperazinamide	319
X L D Y Z		phenylacetyl	morpholinamide	313
X L D F Z		phenylacetyl	4,4-N,N-(dimethyl)-piperazinumamide iodide''	313
X L D F Z		phenylacetyl	amide	313
X L D F P		phenylacetyl	morpholinamide	291
X L D L Z		phenylacetyl	pyrrolidinamide	250
X L D F Z		phenylacetyl	piperidinamide	231
X L D F Z		anilino carbonyl	amide	227
X L D F P		phenylacetyl	amide J = cyclohexyl-Ala'	217

- 34 -

5	X L D F P Z	phenylacetyl	peptide-substituted tetraethylene-pentaamine'	209
	X L D F P	pyridine-3-carbonyl	amide	201
	X L D F Z	phenylacetyl	3,4-dihydroxypyrrrolidinamide	194
	X L D F Z	phenylacetyl	2-(hydroxymethyl)prolinamide	184
	X L D W P	phenylacetyl	amide	184
	R Z F L D F P	B-biotinoyl	2-ε-amidocaproyl amide	150
	X L D V P	phthalimido	amide	140
	Z X F L D F P	ε-amidocaproyl	2-Europium label' amide	118
	X L D V P	phenylacetyl	amide	106
	X L D Z	phenylacetyl	N-(1-(carboxamidomethyl)-caprolactam-3-yl)amide	97
	X L D J P	phenylacetyl	amide J-homo-Phe'	94
	X L D F Z	3,4-dihydroxyphenylacetyl	piperidinamide	81
	X L D F P	benzoyl	amide	75
	X L D V P	pyridine-3-carbonyl	amide	71
	X L D J P	phenylacetyl	amide J-Phenyl-Gly'	61
	X L D F P	3-quinolinecarbonyl	amide	56
	X L D F P	1,2,3,4-tetrahydroquinazoline-2,4-dione-3-yl	amide	56
	X L D Z	phenylacetyl	6,7-fused ring Lactam'	50
	X K D F P	phenylacetyl	amide J-Nle'	49
	P L D F P	free amine	amide	47
	X L D Z	phenylacetyl	N-((1-carboxamidomethyl)-2-oxo-tetrahydroquinolin-3-yl)amide	45
6	X F L D L Z	GlcNAc-O-(CH <sub>2</sub> )-C(O)	piperidinamide	41
	X L D J P	phenylacetyl	amide J = p-nitro-Phe'	40
7	X L D V P	benzoyl	amide	37
	F L D F P	acetyl	amide	35
	X L D F P	benzyloxycarbonyl	amide	34

- 35 -

X L D F Z	(5-phenyl)hydantoinyl	piperidinamide	34
X L D F P	pyrazolecarbonyl	amide	35
F L D F P	acetyl	amide	33
X L D Z	phenylacetyl	N-(2-oxo-tetrahydroquinolin-3-yl)amide	31
X L D Z	phenylacetyl	N-(caprolactam-3-yl)-amide	27
X L D F Z	phenylacetyl	propanolamide	25
X L D Y Z	4-pyridinecarbonyl	piperidinamide	24
X L D Z	phenylacetyl	N-[1-(2-N-morpholinylethyl)-2-oxo-tetrahydroquinoline-3-yl]amide	20
X L D V P	pivaloyl	amide	20
X L D V P	benzoyl	amide	20
X L D Z	phenylacetyl	benzylamide	19
X L D Z	phenylacetyl	phenethylamide	19
X J D F P	benzoyl	amide J-cyclohexyl-Ala'	17
X L D Z	phenylacetyl	N-D-(caprolactam-3-yl)amide	17
X L D S P	phenylacetyl	amide	17
X L D Y Z	phenylacetyl	t-butylester	15
X L D V P	phenylpropionyl	amide	14
X L D Z	phenylacetyl	N-(2-N-morpholinylethyl)amide	13
X J D V P	benzoyl	amide J-cyclohexyl-Ala'	11
F L D V P	free amine	amide	10
X L D F P	2-pyrazinecarbonyl	amide	9
X L D G Z	phenylacetyl	morpholinamide	9
X L D V P	2,3-dimethylbenzoyl	amide	9
X L D V P	3,4-dimethylbenzoyl	amide	8
X L D V P	pyridine-2-carbonyl	amide	8
X L D Z	phenylacetyl	2-N-[1-(N-cyclohexyl)butyrolactam-3-yl]amide	7







- 36 -

	X L D Z	phenylacetyl	N-(1-iso-butyl-2-oxo-tetrahydroquinolin-3-yl)amide	6
	X L D F Z	benzyl	piperidinamide	6
	X L D V P	free amine	amide	5
	X L D V P	cyclohexanecarbonyl	amide	5
	X L D V P	2,6-dimethylbenzoyl	amide	5
	X L D F P	2-quinolinecarbonyl	amide	4
	X L D V P	3-methylvaleroyl	amide	3
	X L D Z	phenylacetyl	N-(tetrahydroisoquinolin-3-yl)amide	3
	P L D F P	free amine	amide	3
	X L D F P	8-quinolinesulfonyl	amide	3
	X L D F Z	phenylacetyl	n-butylamide	3
	X L D V P	4-methylvaleroyl	amide	3
	X L D Y	phenylacetyl	t-butyl ester	3
	X L D Z	phenylacetyl	benzylhydramide	3
	X L D F P	p-bromophenylacetyl	amide	3
	I L D F P	free amine	amide	2
9	X L D F P Z	phenylacetyl	decylamide	2
10	I L D V P I L D V P	free amine	amide	2
	X J D F P	benzylamide	amide J=dicarboxy-Leu <sup>1</sup>	2
	X L D V P	cyclohexanecetyl	amide	2
	X L D Z	phenylacetyl	N'-t-Boc-hydrazide	2
11	I L D F P	free amine	amide	2
	X L D V P	1-naphthoyl	amide	1
	X L D V P	cyclohexanepropionyl	amide	1
	X L D Z	phenylacetyl	N'-benzyl-N'-cyclohexanecarbonylhydrazide	1
	F J D F P	free amine	amide cyclohexyl-Ala <sup>1</sup>	1
	I L D V P	free amine	amide	1

- 37 -

3	I L D V P	free amine	amide	1
12	I J D F P	free amine	amide	1
	X L D V P	cinnamoyl	amide	1
	G P E I L D V P S T	free amine	free acid	1
	X L D F P Z	phenylacetyl	HN(CH <sub>3</sub> ) <sub>2</sub> C(O)NHC <sub>10</sub> H <sub>11</sub>	1
	X L D V P	adamantanecarbonyl	amide	1
	X L D V P	2-naphthoyl	amide	1
13	I L D V P	free amine	amide	1
14	I L D V P	free amine	free acid	1
	X L D F	phenylacetyl	amide	<1
	X L D F Z	phenylacetyl	N-(4-decoyloxy)piperidinamide	<1
	X L D F Z	phenylacetyl	N-(4-stearoyloxy)piperidinamide	<1
	X L D Z	phenylacetyl	hydrazide	<1
15	S F D F S	acetyl	amide	<1
	I J D V P	free amine	amide	<1
	X L D F P	4-bromophenylsulfonyl	J=cyclohexyl-Ala'	<1
	L D F Z	free amine	amide	<1
	J D F Z	J-iso-butyloxycarbonyl	piperidinamide	0
	X L D Z	phenylacetyl	N',N'-dibenzylhydrazide	0
	X L D Z	phenylacetyl	N-(1,3-diphenylprop-2-yl)amide	0
	X L D Z	phenylacetyl	dibenzylamide	0
	X L D F P	phenylsulfonyl	amide	0
	L D V	acetyl	amide	0
	L D F	acetyl	amide	0
	L D F	free amine	amide	0
	L D V	free amine	amide	0
	L L D V P	free amine	free acid	0
	X L D F P	phenylacetyl	amide	0
			scrambled	0

- A lower case letter in bold-faced type is used to designate a D-isomer of the L-amino acid residue designated in single letter code by the same capital letter. Thus, p-D-proline; f-D-phenylalanine; i-D-isoleucine. The N-terminal  $\alpha$ -amine is substituted as shown or indicated to be a "free amine".
- The state of the C-terminal carboxyl of Z as an "amide" ( $-\text{NH}_2$ ) or "free acid" is noted as appropriate.
- Z is an amide formed between the C-terminal pro carboxyl and a tetraethylenepentaamine containing four N-phenylacetyl-LDPP peptides amide-bonded thereto.
- Diethylenetriaminepentaacetatouropium (II) amide-bonded to Z.
- "Homo-Phe" = homophenylalanine; "phenyl-Gly" = phenylglycine; "p-nitro-Phe" = p-nitrophenylalanine;  $\beta$ -carboxy-Asp =  $\beta$ -carboxyaspartic acid; "cyclohexyl-Ala" = cyclohexylalanine; "dicarboxy-Leu" = dicarboxyleucine; and "Nle" = norleucine.
- Relative activities of about one-tenth or less than that exhibited by the peptides of SEQ ID NO:3 are assigned a potency activity of zero.





- 39 -

The data of Table 1 can be used to predict potency in binding inhibition for an unmade or another inhibitor peptide. Thus, one can examine the structures of Table 1 and use the observed relative inhibition potencies to calculate the relative contribution of individual residues or terminal groups. With such calculations in hand, one can then predict the potency of another contemplated peptide within a factor of about 5-10.

For example, comparing the inhibitions exhibited by N-phenylacetyl-Leu-Asp-Val-D-Pro-NH<sub>2</sub> and N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub> one sees that replacement of Val by Phe provides an enhancement of inhibition by a factor of about 3. The differences between the otherwise similar pyridine-3-carbonyl derivatives is similarly about 3.

One could therefore predict that the two N-benzoyl derivatives of the peptides immediately above would similarly be about a factor of three different in their potencies. The data in Table 1 indicate that the observed difference is about a factor of 3.8, thereby verifying the prediction.

Carrying this useful but simplified procedure further, it can be calculated from peptides N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub> and N-phenylacetyl-Leu-Asp-Phe-piperidinamide that D-Pro-NH<sub>2</sub> provides about 1.3-times greater activity than does piperidinamide. Comparing the activity of the latter peptide to that having a morpholinamide group, one calculates an enhancement of about 3.6-fold for a morpholineamide as compared to piperidinamide group. Comparing the peptides N-phenylacetyl-Leu-Asp-Tyr-morpholinamide and N-phenylacetyl-Leu-Asp-Tyr-D-Pro-NH<sub>2</sub>, one finds an enhancement in activity of about a 2.05-fold for morpholinamide over D-Pro-NH<sub>2</sub>. Using those two

- 40 -

calculations, one can then calculate that a morpholinamide group provides about 7.4-times the enhancement of a piperidinamide group ( $2.05 \times 3.6$ ). That predicted enhancement is about 5.7-times greater than that found with the first two peptides ( $7.4/1.3$ ), but within the range of the before-stated predictability.

The data of Table 1 also illustrate the unexpected enhancements in binding inhibition exhibited by an inhibitor peptide contemplated herein as compared to other peptides of the art. For example, the Leu-Asp-Val (LDV) peptide stated as being a minimal peptide required for binding of the VLA-4 receptor in WO 93/12809 exhibited a relative inhibitory binding activity of about 1 as the N-acetyl C-amide derivative and zero as the N-free amine C-amide as compared to the peptide of SEQ ID NO:3. Similar results were observed with Leu-Asp-Phe (LDF) that is not disclosed in that published application, but is an important core peptide here. Contrarily, when an X and Z group as defined herein are added to the termini of either peptide, enhancements in binding inhibition of about 1 to about 3 orders of magnitude were observed.

The data shown in Figs. 1, 2 and 3 also illustrate the unexpected binding inhibitions exhibited by contemplated peptides relative to other peptides of the art. The peptide sequences are shown using single letter code.

For example, Fig. 1 illustrates results of relative in vitro binding inhibition studies carried out using the CS-1 (SEQ ID NO:1) peptide, the CS-1 peptide B12 portion (CS-1 B12; SEQ ID NO:2), the 10-mer peptide used as a standard above, elsewhere herein and in the art (SEQ ID NO:3), and several deletion analogues of the B12 peptide, each containing the Leu-Asp sequence. N-Terminal deletion analogues are shown to the left of

- 41 -

the standard 10-mer, whereas C-terminal deletions are shown to the right of the 10-mer. As is seen, the CS-1 peptide is about three times more potent an inhibitor than is B12, the 10-mer or a 9-mer deletion analogue of the 10-mer. Those latter three peptides were all more potent than the other B12-related peptides.

The similarly obtained data of Fig. 2 illustrate binding inhibition results obtained using deletion analogues of the standard 10-mer peptide.

Here, deletions made at both N- and C-termini are shown to the left of the standard 10-mer to isolate the Leu-Asp-Val sequence at the C-terminus, whereas those shown to the right of the standard 10-mer isolate the Asp-Val-Pro sequence. These peptides and those of Fig. 1 had free N-terminal amine groups and C-terminal carboxyl groups.

The data of Fig. 3 were similarly obtained, but are shown on a log scale so that all of the data could be accommodated. The data of Fig. 3 are shown in five groups, from left to right.

The first group show data for CS-1 peptide and the 10-mer standard. The next three bars shown data for a pentamer C-amide having the sequence including Leu-Asp-Val of the native CS-1 peptide, the enhanced effect of using D-proline instead of the native L-proline, and then the enhancement by use of phenylalanine and D-proline in place of valine and D-proline. The next two bars illustrate the further enhancement obtained over the three previous peptides obtained when a cyclic ring-containing X group, here phenylalanine as the free amine, was used to replace the isoleucine of the native sequence. The fourth group of bars illustrates the effects of three X groups of formula I as compared to the phenylalanine group, using the better peptide sequence of the two adjacent sequence [XLDFp-NH<sub>2</sub>].

- 42 -

Phenylacetyl ( $\phi$ Ac) was used as an X group in the last three peptides where the D-proline Z group of formula I was varied using three cyclic amines (NCy<sup>1</sup>). As is seen, use of a morpholinamide group as Z, along with phenylacetyl as X and phenylalanine as Xaa of formula I, provided the greatest potency in these studies.

Thus, the data of Fig. 3 show inhibitory potencies spanning about three orders of magnitude from the standard 10-mer and peptides of the art, through contemplated peptides that exhibit about a 10-fold enhancement in potency over that standard to those contemplated peptides exhibiting about a 50-fold to about 100-fold enhancement in potency and those exhibiting an enhancement in potency of up to about 1000-fold.

In addition to being more potent than the CS-1 or standard 10-mer peptides, a contemplated inhibitor peptide, particularly a peptide with non-naturally occurring terminal groups such as N-phenylacetyl and C-morpholinamide or D-Pro-NH<sub>2</sub>, is relatively more stable in serum than is the CS-1 peptide. Thus, the inhibitor peptides N-phenylacetyl-Leu-Asp-Phe-morpholinamide and N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub> exhibited no loss of potency after 24 hours in PBS at 7.2-7.4 that also contained 10 percent mouse or human serum. Contrarily, the CS-1 peptide lost its potency in less than one hour under the same conditions.

#### B. Syntheses

The contemplated inhibitors are peptides or peptide derivatives, and as such, can be readily synthesized using well known synthetic methods. Specific synthetic examples are provided hereinafter.

Solid phase synthesis was used for those materials having a C-terminal amino acid amide or free acid residue. Thus, the N-protected, C-terminal residue

- 43 -

was linked to a solid support having a benzhydrylamine substituent. Fmoc amine blocking groups were used in these syntheses, although t-Boc, CBZ or other blocking groups can also be used with other solid supports. Upon  
5 deblocking the Fmoc group with piperidine, another residue was coupled. That coupling was followed by further deblocking, coupling, deblocking etc. steps until a solid phase-linked peptide of desired sequence was prepared. As appropriate to each peptide, an  
10 N-terminal X group was added after a final N-deblocking step or sometimes pre-coupled to the N-terminal residue. The desired peptide and any accompanying functional group protecting groups were removed from the solid support by reaction with trifluoroacetic acid (TFA).  
15 This procedure results in a C-amide-terminated peptide when a benzhydrylamine solid support is used.

Contemplated peptides can also be prepared using t-Boc N-protecting groups and another solid support, or a benzylamino-substituted solid support to  
20 which a p-hydroxymethylphenylcarboxyl (PAM) group is first reacted with the amine of the support to form a carboxamide. The hydroxyl group is then used to form an ester link to the first peptide and standard t-Boc synthetic technology is thereafter followed. Reaction  
25 of the completed, deprotected solid phase-linked peptide with ammonia provides the C-terminal amide peptide discussed before, whereas reaction with another amine such as morpholine or piperidine or other NCy<sup>1</sup> or NCy<sup>2</sup> amine provides a peptide whose C-terminal residue is  
30 amide-bonded to an NCy<sup>1</sup> or NCy<sup>2</sup> group. Reaction of a deprotected, PAM-linked peptide with hydroxide provides the corresponding C-terminal carboxyl group.

In other embodiments, liquid phase peptide syntheses were utilized. For example, morpholine or  
35 other NCy<sup>1</sup> or NCy<sup>2</sup> group was coupled in solution to a

- 44 -

contemplated C-terminal, t-Boc-protected residue using a carbodiimide. The t-Boc protecting group was removed with acid, a further t-Boc-protected residue added, followed by deblockings and further additions. The N-terminal X group such as phenylacetic acid was added after the last t-Boc removal step and the synthesis was completed, except for deprotecting the Asp residue. That step was carried out by catalytic hydrogenation where a benzyl ester protecting group was used.

Regardless of the synthetic method used, an inhibitor peptide is typically recovered and purified prior to use. Recovery and purification techniques are well known and will not be dealt with here.

#### C. Compositions and Process

As noted elsewhere, immune system leukocyte effector or inflammatory cells such as monocytes, T cells and eosinophils bear the VLA-4 receptor on their cell surfaces. Those cells bind to the CS-1 portion of fibronectin present on the surfaces of vascular endothelial cells at an early step in inflammatory cell emigration (trafficking) from the blood in the tissues. These inflammatory cells immunoreact with monoclonal antibody P4C2 discussed in Wayner et al., J. Cell. Biol., 109:1321-1330 (1989), Wayner WO 98/12809, Hemler et al, J. Biol. Chem., 262(24):11478-11485 (1987) and monoclonal antibody HP1/2 of Lobb WO 93/13798 published July 22, 1993.

Once in the tissues, the inflammatory cells enhance the inflammatory response through one or more of several mechanisms. In one mechanism, cytokines and chemoattractants reactants such as interleukin-1 $\beta$  (IL-1 $\beta$ ), IL-2, tumor necrosis factor  $\alpha$  (TNF $\alpha$ ) and lymphocyte-derived chemotactic factor are released by the inflammatory cells and cause further inflammatory cells to emigrate to the area. In another mechanism,

- 45 -

the inflammatory cells mis-recognize cells of the mammal with the inflammatory disease state as being non-self and attack those cells, killing them. These and other mechanisms of immunoinflammatory response enhancement are well known to skilled workers and need not be further elaborated upon here. The fibronectin CS-1 peptide thus mediates inflammatory disease states by assisting emigration of inflammation-enhancing effector cells from the blood into the tissues.

A contemplated inhibitor peptide blocks binding between CS-1 and VLA-4, and inhibits the resulting emigration of inflammatory cells bearing VLA-4 receptors into the tissues, and the exacerbation of the inflammatory condition that results. That inhibition of emigration of inflammatory cells results in a reduction of the fibronectin CS-1/VLA-4-mediated inflammatory response caused by those inflammatory cells, and thereby reduces the observed inflammation.

Particular inflammatory disease states that are mediated by CS-1 and VLA-4, and in which a contemplated inhibitor peptide can diminish inflammation are quite broad. Illustrative of those types of inflammation are asthma, arthritic conditions such as rheumatoid arthritis and osteoarthritis, allograft rejection, various types of skin inflammation, and demyelinating diseases of the central nervous system.

Specific pathological inflammatory conditions in which expression of CS-1 has been found to be implicated and where no such expression is observed in absence of a pathological condition (i.e., in normal tissue) include: rheumatoid arthritis (synovium), osteoarthritis (synovium), skin psoriasis, kidney transplant, asthmatic lung, and lymph node high endothelial venules (HEV) in humans, as well as in the gut of monkeys infected with SIV and those having

- 46 -

inflammatory bowel disease, rabbits having asthmatic lungs and heart transplants, mouse brain in experimental autoimmune encephalomyelitis (EAE) and skin in delayed type hypersensitivity (DTH), and the joints of rats with induced arthritis.

A pharmaceutical composition containing a contemplated inhibitor peptide dissolved or dispersed in a pharmaceutically acceptable carrier or diluent that is preferably aqueous is also contemplated for use in treating a CS-1/VLA-4-mediated inflammatory disease state such as those discussed before. Such a composition contains a CS-1/VLA-4 binding-inhibiting (an inflammation-reducing) amount of a before-discussed, contemplated inhibitor peptide.

Thus, the present invention also contemplates a pharmaceutical composition that can be used in treating one or more of the aforementioned conditions. A contemplated pharmaceutical composition is comprised of a before-described inhibitor peptide that inhibits the binding interaction between VLA-4-containing leukocytes and the fibronectin peptide CS-1 portion expressed on endothelial cell surfaces, which peptide is dissolved or dispersed in a pharmaceutically acceptable diluent in a binding inhibitory (inflammation-reducing) amount. A contemplated pharmaceutical composition is suitable for use in a variety of drug delivery systems. For a brief review of present methods for drug delivery, see, Langer, Science, 249:1527-1533 (1990).

For a contemplated pharmaceutical composition, the dose of the peptide varies according to, e.g., the particular peptide, the manner of administration, the particular disease being treated and its severity, the overall health and condition of the patient, and the judgment of the prescribing physician or veterinarian. A pharmaceutical composition is intended for parenteral,



- 47 -

topical, oral or local administration, such as by aerosol or transdermally, for prophylactic and/or therapeutic treatment. A pharmaceutical composition can be administered in a variety of unit dosage forms depending upon the method of administration. For example, unit dosage forms suitable for oral administration include powder, tablets, pills, capsules and dragees.

Preferably, a pharmaceutical composition is administered intravenously. Thus, a composition for intravenous administration is particularly contemplated that comprises a solution of a contemplated inhibitor peptide dissolved or dispersed in a pharmaceutically acceptable diluent (carrier), preferably an aqueous carrier. A variety of aqueous carriers can be used, e.g., water, buffered water, 0.9 percent saline, buffered aqueous ethanol solutions and the like. These compositions can be sterilized by conventional, well known sterilization techniques, or can be sterile filtered. The resulting aqueous solutions can be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile aqueous solution prior to administration. A composition can contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, etc.

The concentration of inhibitor peptide utilized is usually at or at least about 0.0001 percent to as much as about 0.1 percent by weight and is selected primarily by fluid volumes, viscosities, etc.,

- 48 -

in accordance with the particular mode of administration selected.

Thus, a typical pharmaceutical composition for intravenous infusion can be made up to contain 250 ml of sterile Ringer's solution normal saline or PBS, and about 0.25 mg to about 25 mg of the inhibitor peptide. Actual methods for preparing parenterally administrable compounds are known or apparent to those skilled in the art and are described in more detail in for example, Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, PA (1985), which is incorporated herein by reference.

For solid compositions, conventional nontoxic solid diluents (carriers) may be used which include, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin, talcum, cellulose, glucose, sucrose, magnesium carbonate, and the like. For oral administration, a pharmaceutically acceptable nontoxic composition is formed by incorporating any of the normally employed excipients, such as those carriers previously listed, and generally 10-95 percent of active ingredient, that is, a before-described inhibitor peptide preferably about 20 percent (see, Remington's, supra), preferably using an enteric coating to pass a solid dose through the stomach and into the intestine.

For aerosol administration, a contemplated inhibitor peptide is preferably supplied in solution such as aqueous ethanol or DMSO solution along with a surfactant and propellant. Typical percentages of an inhibitor peptide are about 0.0001 percent to about 0.1 percent by weight, and preferably about 0.0001 percent to about 0.001 percent. The surfactant must of course, be nontoxic, and preferably soluble in the propellant. Representative of such agents are the esters or partial

- 49 -

esters of fatty acids containing from 6 to 22 carbon atoms, such as caproic, octanoic, lauric, palmitic, stearic, linoleic, linolenic, olesteric and oleic acids with an aliphatic polyhydric alcohol or its cyclic anhydride such as, for example, ethylene glycol, glycerol, erythritol, arabitol, mannitol, sorbitol, the hexitol anhydrides derived from sorbitol, and the polyoxyethylene and polyoxypropylene derivatives of these esters. Mixed esters, such as mixed or natural glycerides can be employed. The surfactant can constitute about 0.1 to about 20 percent by weight of the composition, and preferably about 0.25 to about 5 percent. The balance of the composition is ordinarily propellant. Liquefied propellants are typically gases at ambient conditions, and are condensed under pressure. Among suitable liquefied propellants are the lower alkanes containing up to 5 carbons, such as butane and propane; and preferably fluorinated or fluorochlorinated alkanes. Mixtures of the above can also be employed. In producing the aerosol, a container equipped with a suitable valve is filled with the appropriate propellant, containing the finely divided compounds and surfactant. The ingredients are thus maintained at an elevated pressure until released by action of the valve. A pump-activated spray using air as propellant (atomizer or nebulizer) is also contemplated.

For example, for the treatment of asthma in rabbits, the dose of a contemplated peptide is in the range of about 1 to 100 mg/day for a 2-3 kg animal. For a human asthma patient, that dose is in the range of about 1 to about 100 mg/day for a 70 kg patient. Administration for asthma is typically by aerosol from a nebulizer. Ideally, therapeutic administration should begin as soon as possible after the attack begins.

- 50 -

A pharmaceutical composition containing an inhibitor peptide can be administered for prophylactic and/or therapeutic treatments. In therapeutic applications, a composition is administered to a patient already suffering from a disease, as described above, in an amount sufficient to inhibit binding between VLA-4-expressing leukocytes and endothelial cells that express the CS-1 peptide portion; i.e., reduce inflammation and thereby at least partially arrest the symptoms of the disease and its complications. An amount adequate to accomplish this is defined as "therapeutically effective dose", or a "binding-inhibiting amount" or an "inflammation-reducing amount". Amounts effective for this use depend on the severity of the disease and the weight and general state of the patient, but generally range from about 1 mg/kg to about 500 mg/kg of inhibitor peptide per day, with dosages of from about 1 mg/kg to about 10 mg/kg of a compound per day being more commonly used.

In prophylactic applications, a composition containing a contemplated peptide is administered to a patient susceptible to or otherwise at risk of a particular disease. Such an amount is defined to be a "prophylactically effective dose" and is also an amount sufficient to inhibit binding of VLA-4-expressing leukocytes to CS-1 peptide-expressing endothelial cells. In this use, the precise amounts again depend on the patient's state of health and weight, but generally range from about 1 mg/kg/day to about 500 mg/kg/day, more commonly from about 1 mg/kg/day to about 20 mg/kg/day.

Another way to assess a binding-inhibiting amount of a contemplated inhibitor peptide is to compare binding inhibition exhibited by the peptide to that provided by CS-1 or the 10-mer standard in an in vitro

- 51 -

study. One convenient way to make that comparison is by use of  $IC_{50}$  values of the two compared materials, and base the amount used on the amount of CS-1 or standard 10-mer peptide and an amount of the inhibitor peptide that is a multiple of the  $IC_{50}$  value for that reference compound.

Typically, a compound whose  $IC_{50}$  value is at least about one-tenth that of the standard 10-mer (ten-times more potent), when used at one-tenth the molar amount of the 10-mer standard is a useful binding-inhibiting amount. More preferably, the amount is about one-fiftieth the amount of the 10-mer. More preferably still, the amount is equal to about one-hundredth that of the 10-mer. Inasmuch as those amounts inhibit binding by about 50 percent, greater concentrations that inhibit binding still further are preferred.

Thus, for in vitro use, a minimal CS-1/VLA-4-inhibiting amount is the  $IC_{50}$  value. For in vivo use, the CS-1/VLA-4-inhibiting amount usually used begins with the  $IC_{50}$  value concentration, and can decrease as required or one can increase to the solubility limit of the peptide in the utilized aqueous medium; i.e., the aqueous medium at pH 7.2-7.4 used such as normal saline where parenteral administration is used or intestinal fluid where oral administration is used.

Single or multiple administrations of a composition can be carried out with dose levels and pattern being selected by the treating physician or veterinarian. In any event, a pharmaceutical composition is formulated to provide a quantity of an inhibitor peptide sufficient to effectively treat the patient.

A process for treating fibronectin CS-1/VLA-4-mediated inflammation is also contemplated. In accordance with such a process, a before-described,

- 52 -

contemplated inhibitor peptide is administered to a mammal in need of such a treatment such as a mammal having such inflammation or prophylactically as prior to an allograft. This administration is preferably via a before-discussed pharmaceutical composition. The peptide is administered in an inflammation-reducing (CS-1/VLA-4 binding inhibiting) amount. The mammal such as mouse, rat, rabbit, monkey or human is maintained until the peptide is eliminated by a natural bodily process. Multiple administrations in a single day, over a period of days or weeks, or for the life of the host mammal, where the mammal is the recipient of an allograft, are contemplated, as are single administrations.

Methods for determining an amount sufficient to inhibit binding between CS-1 and VLA-4 have already been discussed, particularly for in vitro studies. For in vivo uses, there are many published assays to determine if inflammation has been reduced by a particular treatment. For example, one can assess the number of painful joints in an arthritic patient or the patient's mobility before and after treatment. Reduction of effects of an asthma attack can be assayed by measurement of dynamic compliance or lung resistance in laboratory animals as is also well known. The amount of edema observed in DTH is also readily measurable, as are the effects of allograft rejection or its absence compared to standard controls.

### Best Mode for Carrying out the Invention

#### Example 1: Synthesis of X-Leu-Asp-Phe-Z

Protected amino acids, Boc-Phe-OH, Boc-Asp(OBn)-OH and Boc-Leu-OH were purchased from NOVA Biochem Co., La Jolla, CA and were used without further purification. Phenylacetic acid, morpholine,

- 53 -

diisopropylethylamine (DIEA) and 1-hydroxybenztriazole (HOBt) were obtained from Aldrich Chemical Co., Milwaukee, WI. Ethyl-3-(3-dimethylamino)-propylcarbodiimide•HCl (EDC) was obtained from Bachem Co., Torrance, CA. 4 Normal HCl in dioxane was obtained from Pierce Co., Rockford, IL used as received.

The <sup>1</sup>HNMR spectra were recorded on a GE QE-300, 300 MHz NMR spectrometer.

A. Preparation of Boc-Phe-Morpholinamide

A 250 ml flask was charged with Boc-Phe-OH (10 g, 38 mmol), morpholine (3.3 g, 38 mmol) and 1-hydroxybenztriazole (5.1 g, 38 mmol) in 100 ml dry dimethylformamide (DMF). To this solution was added diisopropylethylamine (DIEA) at zero degrees C until the pH value reached 8, followed by addition of EDC (8.8 g, 46 mmol). The solution was slowly warmed to room temperature. The mixture was stirred for eight hours at room temperature (about 22°C). The DMF was removed by vacuum evaporator, ethyl acetate and water were added, and the layers were separated. The aqueous layer was extracted with ethyl acetate (50 ml x 2), combined extracts were washed with 1N HCl, saturated NaHCO<sub>3</sub>, water and brine, dried with MgSO<sub>4</sub>, filtered and concentrated to give a colorless liquid (12.8 g, 37.7 mmol; 99 percent yield) that was characterized by <sup>1</sup>HNMR as Boc-Phe-morpholine.

B. Preparation of HCl-Phe-Morpholinamide Salt

Boc-Phe-morpholinamide (12.8 g, 38 mmol) was placed in a 250 ml flask, then 4N HCl in dioxane (30 ml) was added. The mixture was stirred for six hours at which time TLC (silica gel; CHCl<sub>3</sub>:MeOH:acetic acid, 90:8:2) indicated that the reaction was completed. Dioxane and excess HCl were removed. A white solid, identified by <sup>1</sup>HNMR as HCl-Phe-morpholinamide, was obtained in 100 percent yield (10.3g, 38 mmol).

- 54 -

C. Preparation of N-Phenylacetyl-Leu-Asp( $\beta$ -O-Benzyl)-Phe-Morpholinamide

Boc-Asp( $\beta$ -OBn)-OH, Boc-Leu-OH and phenylacetic acid were sequentially added to HCl-Phe-morpholinamide using the coupling and deprotection procedures, described above. The white solid, thus obtained, was crystallized from ethyl acetate and hexane, and identified by  $^1\text{H}$ NMR as the desired ester in 95 percent yield.

D. Synthesis of N-Phenylacetyl-Leu-Asp( $\beta$ -O-Bn)-Phe-Morpholinamide

To a solution of the above benzyl ester (10 g, 15 mmol) in methanol (100 ml) was added 10 percent palladium-charcoal (2.0 g). The flask containing this mixture was evacuated and then filled with hydrogen three times. The mixture was then vigorously stirred under a hydrogen atmosphere about five hours until the hydrogenolysis was complete, as indicated by TLC ( $\text{CHCl}_3$ :MeOH:acetic acid, 90:8:2). The reaction mixture was filtered through celite, and the methanol was removed, affording a white solid that was characterized by  $^1\text{H}$ NMR as XLDFZ (8.4 g, 14.5 mmol) in 97 percent yield.

Peptides having other than carboxamide [C(O)-NH<sub>2</sub>] C-terminal amide-linked Z groups were similarly prepared.

Example 2: Exemplary Solid Phase Peptide Syntheses

Fmoc protected amino acids, hydroxybenzotriazole (HOBt) and Rink amide MBHA resin were obtained from Nova Biochem, La Jolla, CA. Diisopropylcarbodiimide (DIC) was obtained from Chem Impex Inc., Chicago, IL. Piperidine was obtained from Aldrich Chemical Company, St. Louis, MO. Dimethylformamide (DMF), isopropanol (IPA),



- 55 -

dichloromethane (DCM), and dimethylacetamide (DMA) were obtained from Burdick and Jackson, Muskegon, MI. All of the above reagents were used as supplied by the manufacturer, with no further purification.

5           The standard deprotection/coupling cycle iterated during this synthesis is described in terms of the first coupling of Fmoc-Pro to the Rink amide MBHA resin:

10           The Fmoc-MBHA resin (10.6 g., 5 mmoles) was treated with 20 percent piperidine in DMF (130 ml) for three minutes. The solution was removed by filtration and the resin was again treated with 20 percent piperidine in DMF (130 ml) for 17 minutes. The solution was removed by filtration and the resin was washed five  
15           times with DMF (130 ml each), two times with IPA (130 ml) and two times with DMF (130 ml). The HOBt ester of Fmoc-D-proline (formed by reacting a solution of 10 mmoles Fmoc-D-proline and 10 mmoles HOBt in 50 ml DMA with 12 mmoles DIC for 20 minutes at room temperature),  
20           in DMA (50 ml), was added to the resin and allowed to react for two hours. The resin was washed five times with DMF (130 ml) and two times with DCM (130 ml). The coupling of amino acid to the resin was checked by standard Kaiser's test.

25           The above cycle was iterated for each of the subsequent amino acids: Fmoc-Phe, Fmoc-Asp( $\beta$ -OBn), Fmoc-Leu, and phenylacetic acid. The resin was dried in vacuo for 24 hours and then allowed to react with 95 percent TFA/5 percent H<sub>2</sub>O (60 ml) for two hours at room  
30           temperature. The TFA solution of the peptide was separated from the resin by filtration and the TFA was vacuum evaporated. The solid residue was crystallized from anhydrous ethanol to yield 1.8 g of product, N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub>. The peptide was

- 56 -

characterized by amino acid analysis on HP Amino Quant 1090 and NMR, and the purity of the peptide was checked by HPLC (WATER HPLC Systems).

5      Example 3:      In Vitro Binding Assays

        Jurkat cells (ATCC TIB 152), a human T lymphoblastic line, labeled with <sup>51</sup>chromium were used to assay in vitro binding inhibition provided by various peptides discussed herein. Costar™ 96 well flat-bottom microtiter plates (catalog No. 9050, Cambridge, MA) were found to provide the best results in these assays.

        The plates were prepared as follows: The 25-mer CS-1 peptide (SEQ ID NO:1) dissolved at 0.5-1 µg/ml in a buffer of 0.1M NaHCO<sub>3</sub> at pH 9.5 that also contained 10 µg/ml of bovine serum albumin (BSA) or a conjugate of the CS-1 peptide linked to ovalbumin (CS-1-OVA) dissolved at 1-2.5 µg/ml in the same buffer was used as the substrate. Each well of the microtiter plates was coated with 50 µl of substrate or buffer alone for controls. The wells were permitted to dry out completely and were then rinsed twice with PBS at pH 7.4. Non-specific binding sites of each well were then blocked using 200 µl per well of RPMI/1 percent BSA for two hours at room temperature. Both solid phase-affixed substrates provided similar results.

        Jurkat cells (3-5X10<sup>6</sup> cells) were placed into a 15 ml Falcon™ round-bottom tube with a cap. The tube was centrifuged, and the extra medium was then removed.

        Two hundred microliters of a <sup>51</sup>Cr labeling solution were added to the centrifuged cells and maintained in contact with the cells for 90-120 minutes in a warm room. This procedure typically provides about 50,000-100,000 cpm/well with about 80-100 percent cell viability. Longer contact times provide a greater amount of labeling but lower cell viability.

- 57 -

The labeled cells are washed with (i) complete medium, (ii) 1mM EDTA/PBS and then (iii) RPMI/1 percent BSA free of serum components. The cells are centrifuged after each washing. The cells are finally resuspended in serum-free RPMI/1 percent BSA at a concentration of  $1 \times 10^6$  viable cells/ml, which provides a concentration that is diluted by one-half in the assay.

Inhibitor peptides are prepared as stock solutions at 20 mg/ml in DMSO in 1.5 ml cryogenic screwcap vials, and were stored at  $-70^\circ\text{C}$ . Using Flow™ round-bottom or V-bottom microtiter plates, the inhibitor peptides were prepared at twice the assay concentration in RPMI/1 percent BSA at 60  $\mu\text{l}$ /well.

Four initial dilutions were typically used. For less potent peptides such as the standard 10-mer of SEQ ID NO:3, the initial dilutions were 500  $\mu\text{g}/\text{ml}$ , 100  $\mu\text{g}/\text{ml}$ , 20  $\mu\text{g}/\text{ml}$  and 4  $\mu\text{g}/\text{ml}$ . For more potent peptides such as N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub>, the typical initial concentrations were 10  $\mu\text{g}/\text{ml}$ , 2  $\mu\text{g}/\text{ml}$ , 0.4  $\mu\text{g}/\text{ml}$  and 0.08  $\mu\text{g}/\text{ml}$ .

The  $^{51}\text{Cr}$ -labeled cells ( $1 \times 10^6$  cells at 60  $\mu\text{l}$ /well) were then admixed with the diluted peptide solutions. The admixtures were maintained at room temperature (about  $22^\circ\text{C}$ ) for 30 minutes.

One hundred microliters of each inhibitor peptide/cell admixture were transferred to the substrate-coated wells. This was done in triplicate for each dilution. The resulting plates were incubated for 30 minutes at  $37^\circ\text{C}$  and then washed gently three times with RPMI/1 percent BSA at 200  $\mu\text{l}$ /well. Binding was observed microscopically, particularly after the second wash.

The bound cells were then lysed by the addition of a 0.5 percent solution of sodium dodecylsulfate in water at 100  $\mu\text{l}$ /well. The resulting

- 58 -

solutions were then processed for counting and calculation of  $IC_{50}$  values following usual procedures. Appropriate positive and negative controls were used with each plate so that the results of separate assays could be normalized and compared.

The data of Table 1 are so normalized. The absolute  $IC_{50}$  value for the peptide N-phenylacetyl-Leu-Asp-Phe-morpholinamide is 0.18-0.30  $\mu M$ .

10      Example 4:      Delayed Type Hypersensitivity in Mice

                 A.    Initial Study

                 Chisholm et al., Eur. J. Immunol., 23:682-688 (1993) reported in vivo results of blocking VLA-4 interactions in a murine contact hypersensitivity model using rat anti-mouse  $\alpha$ -4-antibodies. Those workers noted that therapeutics designed to interfere with VLA-4-mediated adhesion should be effective blockers of in vivo inflammatory processes.

                 An adoptive transfer delayed-type hypersensitivity murine model has been developed using splenic T cells primed to oxazolone. This model is described in Elices et al., Clin. Exp. Rheum., 11(Suppl. 8):577-580 (1993), whose procedures were followed here.

                 Thus, BALB/c mice were shaved on the belly and painted (50  $\mu l$  on the belly and 5  $\mu l$  on each paw) with three percent oxazolone in acetone/olive oil (4:1) at days zero and 1. At day 5, the mice were sacrificed, their spleens removed, and splenic T cells were obtained via nylon wool columns.

                 Normal saline or saline containing  $25 \times 10^6$ /animal of the oxazolone-immune T cells were separately injected into naive mice. The mice were then challenged by painting 10  $\mu l$  of 2 percent oxazolone onto one ear each. All procedures were carried out under sterile conditions and in endotoxing-free buffers.

- 59 -

Prior to challenge or saline injection, the mice were implanted with pumps that subcutaneously administered normal saline, normal saline containing the peptide N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub>, or normal saline containing a peptide with a scrambled sequence continually at 6 mg/kg/day for a 24-hour time period. The swelling diameter at the site of challenge or saline injection was measured with a microcaliper 24 hours thereafter.

The results of this study are shown in the graph of Fig. 5 for the saline and recited inhibitor peptide treatments. Although there is a slight overlap in the data possibly due to non-CS-1-mediated inflammation, it is clear that administration of a contemplated inhibitor peptide reduced this type of CS-1/VLA-4-mediated immunoinflammation as compared to the untreated controls. Use of the scrambled peptide provided no reduction of inflammation. Another study using 600 mg/kg/day of the same peptide showed no toxicity caused by the peptide.

#### B. Expanded Study

An expanded study was carried out as described above except that a scrambled peptide control was not used using further inhibitor peptides of Table 1 with larger groups of mice. Statistical analyses were carried out versus injection of vehicle alone. Those results are shown below in Table 2.

- 60 -

Table 2\*

	<u>Peptide</u>	<u>Percent Inhibition</u>	<u>p Values</u>	<u>n</u>
5	$\phi$ Ac-Leu-Asp-Phe-D-Pro-NH <sub>2</sub>	36	0.0002	30
	$\phi$ Ac-Leu-Asp-Phe	29	0.015	8
	$\phi$ Ac-Leu-Asp-Phe	30	<0.0001	24
10	Phth-Leu-Asp-Val-D-Pro-NH <sub>2</sub>	--	N.S.	8
	$\phi$ Ac-Leu-Asp-Phe	--	N.S.	22

15

\*  $\phi$ Ac = N-phenylacetyl;  
 Phth = N-phthalimido; and  
 N.S. = not significantly different from vehicle.

20

25

30

A post-challenge reduction in swelling was noted for four of the five inhibitor peptides studied, with three of the inhibitor peptides exhibiting a statistically significant reduction in swelling. The fourth peptide that exhibited inhibition did so reproducibly, but not at a statistically significant level. That fourth compound is very hygroscopic. It is therefore believed that substantially less than the stated 6 mg/kg/day dose was actually administered, because of that hygroscopicity, and that that lowered dose was responsible for the lack of a statistically significant result.

35

40

The transferred T cell is the effector cell in the adoptive immune response examined here. T cells express both VLA-4 and VLA-5 that interact with the CS-1- and RGD-containing portions of fibronectin, respectively, during an inflammatory immune response. Ferguson et al., Proc. Natl. Acad. Sci., USA, **88**:8072-8076 (1991) showed that about 50  $\mu$ g/ml of the standard peptide used here (SEQ ID NO:3) when admixed with immune T cells could abrogate a transferred immune response such as the response studied here. Those workers also

- 61 -

taught that separate administration of the peptide and T cells did not lead to that abrogation.

Here, it is seen that separate administration of 6  $\mu$ g/ml of a contemplated inhibitor peptide provided  
5 substantial inhibition of the immune response. It is also seen that the inhibitor peptide and T cells need not be premixed here as they were required to be in the Ferguson et al. results.

The N-phthalimido derivative exhibited an  
10 average swelling that was slightly higher than that exhibited by vehicle alone for a reason that is presently unclear although only one concentration was utilized here and the N-phthalimido derivative exhibited about 1/5 to 1/6 the activity of the  $\phi$ Ac-Leu-Asp-Phe-D-  
15 Pro-NH<sub>2</sub> in the in vitro assay of Table 1 and so it is not particularly unexpected that that less active compound showed no activity at the low end of the activity range.

20 Example 5: Treatment of Asthmatic Rabbits

Six New Zealand white rabbits were immunized with house dust mite antigen from birth through four months of age. Upon immunization, three rabbits received a single nebulizer administration of the  
25 inhibitor peptide N-phenylacetyl-Leu-Asp-Phe-morpholinamide in aqueous 50 percent ethanol as diluent in an amount of 100 mg/kg, and the other three received diluent alone. All of the rabbits were challenged with house dust mite antigen about 15-30 minutes after  
30 administration of the peptide, with those animals not receiving peptide serving as controls.

Once immunized and challenged, the inflammatory state subsides to a basal level within about three weeks. The three animals used as controls  
35 were thereafter used as subjects for receipt of an

- 62 -

inhibitor peptide, and the three rabbits that initially received the peptide can serve as controls.

Such a crossover study was done here. Thus, the three initial control rabbits were treated with the above inhibitor peptide in the above diluent at a time more than three weeks after the above study, and the three previous recipients of the peptide were administered the diluent alone. All six were then challenged again.

Initial pulmonary function, measured by dynamic compliance ( $C_{dyn}$ ) and lung resistance ( $R_L$ ), and bronchoalveolar lavage (BAL) to obtain an effector cell count, here eosinophils, were conducted prior to administration of the peptide or diluent for both portions of this crossover study. Similar assays were then taken one-half hourly after challenge for six hours (early phase allergic reaction) and at 24 hours after challenge (late stage allergic reaction) for both portions of this study.

These studies were conducted by Dr. W. J. Metzger of East Carolina University, Greenville, N.C. as described by W.J. Metzger in CRC Handbook of Late Phase Reactions, W. Dorsch, ed., Chapter 35, CRC Press, Boca Raton, FL (1990) pages 347-362.

The results of this study for the pulmonary function parameters are shown in Fig. 4A and Fig. 4B, in which data for the challenged, inhibitor peptide-treated animals are shown as open circles and data for the challenged, untreated, control animals are shown in blackened circles. These data are averaged values from both portions of the study.

As is seen from Fig. 4A, the  $C_{dyn}$  value for the challenged and treated animals stayed at about the initial value for the whole six hours. The  $C_{dyn}$  for the challenged, untreated animals quickly fell to about 40



- 63 -

percent of the initial value and then stayed at about that value for the whole six hours.

The data of Fig. 4B show that the  $R_L$  values for the challenged, inhibitor peptide-treated animals remained between the initial value and about 200 percent of that value for the whole six hours, with a slight rise near the end of that time period. The  $R_L$  values for the challenged, but untreated animals rose to about 200-300 percent in the first two hours after challenge and rose to about 400-1200 percent for the last four hours.

A summation of the averaged data for the inhibitor-treated, challenged animals compared to the challenged control animals for early (2-4 hours) and late (24 hours) phases of this inflammatory immune response is provided in Table 2, below.

Table 2

IN VIVO Efficacy of  $\phi$ Ac-Leu-Asp-Phe-Morph\*

<u>Parameter</u>	<u>Phase</u>	<u>% Reduction</u>
$C_{dyn}$	Early	94.4
	Late	86.6
$R_L$	Early	80.1
	Late	82.6

---

\* N-Phenylacetyl-Leu-Asp-Phe-morpholinamide

The BAL count from these studies indicated an 88.1 percent reduction in eosinophils after 24 hours in the inhibitor peptide-treated, challenged animals as compared to the untreated, challenged animals in the crossover study.

As can be seen from the above data and those of Figs. 4A and 4B, aerosol administration of an

- 64 -

inflammation-reducing amount of a contemplated peptide greatly reduced the asthmatic response in the treated animals as compared to those receiving no treatments.

5      Example 6:      Rabbit Cardiac Allograft Model

                 New Zealand white rabbit SPF hearts were allografted into the necks of similar rabbits to assay a graft-vs-host immunorejection model and the affect of a contemplated peptide on that immunoinflammatory  
10      response. These studies were carried out by Dr. M. Rabinovitch of the Hospital for Sick Children, Toronto, Ontario, Canada.

Experimental animal model

15                  New Zealand white female rabbits (Charles River Lab., Saint Laurent, Quebec), between 3.5 and 4 kg underwent heterotopic cardiac transplant following an experimental protocol previously described [Alonso et al., Am. J. Pathol., 87:415-442 (1977); Clausell et al.,  
20      Circulation, 89:2768-2779 (1994)] and approved by the Animal Care Committee of The Hospital for Sick Children, Toronto. The animals were unselected to favor an HLA-mismatch, the host rabbits were Pasteurella-free and the donors, outbred animals. Both host and donor rabbits  
25      were fed Purina 5321-0.5 percent cholesterol diet (Research Diets Inc., New Brunswick, NJ), a strategy that has proven useful in accelerating the process of allograft arteriopathy [Alonso et al., Am. J. Pathol., 87:415-442 (1977)]. The diet was commenced four days  
30      prior to the transplant and continued in the recipient of the transplant until the completion of the experimental period.

                 The technique of heterotopic cardiac transplantation has been previously described [Clausell  
35      et al., Circulation, 89:2768-2779 (1994)]. Briefly, a

- 65 -

vertical incision was performed in the anterior aspect of the neck of the recipient rabbit and the left common carotid artery and the ipsilateral external jugular vein were isolated. The cardiac allograft was placed in the neck by anastomosing the aorta end-to-side to the recipient's carotid artery and the pulmonary end-to-side to the recipient's external jugular vein, following a total period of ischemia for the donor hearts of approximately 30 minutes. Postoperative care was in compliance with the Principles of Laboratory Animal Care formulated by the Canadian National Society for Medical Research.

The treatment consisted of an inhibitor tetrapeptide, N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub>, (treated), derived from the leucine-aspartate-valine (LDV) sequence [Komoriya et al., J. Biol. Chem., 266:15075-15079 (1991; Wayner et al. J. Cell Biol., 116:489-497 (1992)], that enhanced inhibition in vitro here, and a scrambled form of the same synthetic tetrapeptide N-phenylacetyl-Asp-Leu-Phe-D-Pro-NH<sub>2</sub>, (control), resulting in no inhibition of VLA-4. Both peptides were synthesized at Cytel Corporation, San Diego, CA.

Beginning the day of the transplant, the animals were randomized and treated with either scrambled peptide (control group) at 1 mg/kg s.c. or the inhibitor peptide at 1 mg/kg s.c. The doses of the peptides were empirically extrapolated to the in vivo model based on preliminary in vitro studies, of Table 1. No other immunosuppression therapy was administered. The grafts were monitored daily by palpation and maintained for 7 to 8 days, a previously described endpoint that was associated with myocardial rejection (impaired cardiac contractility) and development of the allograft arteriopathy in this model [Clausell et al.,

- 66 -

Circulation, 89:2768-2779 (1994)]. A total of fourteen animals were studied in the control (n=7) and treated (n=7) groups.

5     Preparation of the hearts

          The animals were sacrificed using a lethal dose of euthanol (480 mg i.v.) (MTC Pharmaceutical, Cambridge, ON), host and donor hearts were removed, and the coronary arteries were perfused with saline through  
10     the aorta, followed by light fixation by perfusion with 2 percent paraformaldehyde (Sigma, St, Louis, MO). Because of previous descriptions indicating that cardiac allograft arteriopathy in the rabbit was equally distributed throughout the coronary circulation [Foegh  
15     et al., Transplant Proc., 21:3674-3676 (1989)] and those of Dr. Rabinovitch and co-workers [Clausell et al., Circulation, 89:2668-2779 (1994)], the hearts were sectioned transversally from base to apex. Different  
20     sections of the hearts were either saved in 10 percent formalin (BDH Inc., Toronto, ON) for light microscopy studies or immediately frozen in O.T.C. Compound Tissue Tek (Miles Inc., Elkart, IN) for specific immunohistochemistry studies.

25     Grading of rejection

          Tissue specimens from the donor hearts were stained with hematoxylin:eosin for histological grading of rejection according to a modified Billingham's criteria [Billingham, Hum. Pathol., 10:367-386 (1979)].  
30     The sections were graded by one of the senior pathologists (Dr. G.J. Wilson) at the Hospital for Sick Children without knowledge of whether the donor hearts came from control or inhibitor peptide treated animals.

- 67 -

Quantitative assessment of host and donor coronary  
arteries by light microscopy

Three different paraffin-embedded tissue  
5 sections from host and donor hearts from both control  
and treated rabbits were stained by the Movat  
pentachrome method for light microscopy. Morphometric  
analysis was performed using a Zeiss microscope attached  
to a computer-generated video analysis system  
10 (Perceptics Inc., NuVision software), as described in  
Clausell et al., Circulation, 89:2768-2779 (1994). The  
number of vessels with intimal lesions were counted in  
all three heart sections from each animal studied and  
are expressed as a percentage of the total vessel  
15 number.

In the host hearts, 999 vessels in the control  
group and 1054 vessels in the treated group were  
analyzed. In the donor hearts, 827 vessels in the  
control group and 617 vessels in the treated group were  
20 analyzed. To determine the severity of intimal  
thickening, the diameter of each traceable vessel in all  
three sections was measured and the coronary arteries  
were categorized as small (diameter  $<100\mu\text{m}$ ), medium  
(diameter  $>100<500\mu\text{m}$ ) and large (diameter  $>500\mu\text{m}$ ). The  
25 degree of intimal thickening was then quantitatively  
assessed in each vessel size category as previously  
described in Eich et al., Circulation, 87:261-269  
(1993). The areas encompassed by the outer medial layer  
(ML), the internal elastic lamina (IEL) and lumen were  
30 measured in each affected vessel, and the area of  
intimal thickening (IT) related to the vessel area was  
calculated by the formula  $\text{IT} = \text{IEL-lumen area} / \text{ML-lumen}$   
area X 100.

- 68 -

Immunohistochemistry studies

In all immunohistochemistry analyses, coronary arteries from host and donor hearts were compared, with and without intimal thickening in the different size ranges from control and treated groups. The relative abundance of each specific antigen studied in the sections examined was graded semi-quantitatively as minimal (+/-), little (+), moderately abundant (++) to very abundant (+++) by two qualified investigators in the Rabinovitch laboratory. The final scoring was based on individual gradings that reached 90 percent agreement.

(1) Characterization of inflammatory cells

To characterize the presence of an immune-inflammatory reaction in the allograft coronary arteries in both groups studied, immunoperoxidase staining was performed using monoclonal antibodies to rabbit MHC Class II antigens and rabbit T cells (a kind gift from Dr. Peter Libby, Brigham and Women's Hospital, Boston, MA) and also to rabbit macrophages (RAM 11, Dako Corp., Carpinteria, CA). The sections were air-dried for two hours, fixed in acetone for 20 minutes, and rinsed with D-PBS (Gibco, Burlington, ON) /0.1 percent BSA (Boehringer-Mannheim, Mannheim, Germany). Endogenous peroxidase activity was blocked by immersing the sections in PBS/0.1 percent BSA + 3 percent hydrogen peroxide (BDH) for 30 minutes. After a non-specific blocking step using 10 percent normal goat serum (Sigma), the antibodies were applied to the sections for 1 hour at a 1:10 dilution at room temperature. The sections were then rinsed, incubated with goat anti-mouse peroxidase-conjugated secondary antibody (Bio-Rad, Richmond, CA) at a 1:50 dilution at room temperature for 45 minutes and developed with 3,3'-diaminobenzidine

- 69 -

(DAB) (Sigma) for 10 minutes. Control sections were treated with normal mouse isotypic IgG (Dako Corp.).

5           (2)   Immune-detection of cellular adhesion  
              molecules

10           To assess the influence of treatment on the expression of adhesion molecules in allograft coronary arteries, immunoperoxidase staining for ICAM-1 and VCAM-1 was performed on frozen sections of both host and donor hearts from the control and the treated groups. Monoclonal antibodies to ICAM-1 (mAb Rb2/3) and to VCAM-1 (mAb Rb1/9) were kindly supplied by Dr. Myron Cybulsky (Brigham and Women's Hospital, Boston, MA) and were used  
15           at a concentration of 1:10 for 1 hour at room temperature. The procedure for immunostaining was essentially the same as described above.

20           (3)   Assessment of fibronectin  
              Fibronectin expression in coronary arteries of host and donor hearts from both control and treated groups was determined by performing immunoperoxidase staining using frozen sections. A monoclonal antibody anti-cellular fibronectin (Chemicon Int. Inc., Temecula,  
25           CA) was used at a dilution of 1:100 for 1 hour at room temperature and the remaining details of the immunohistochemical procedure are essentially the same as described above. This antibody does not recognize plasma fibronectin.

30           Statistical Analysis

              The data were expressed as mean +/-SE. In analyses related to the incidence and severity of lesions from both control and treated groups, the  
35           Student's  $t$  test was used to test significance. The correlation among categorical variables from the

immunohistochemistry studies, considered positive if  $> +$  in the two groups (control and treated), was analyzed using Fisher's exact test. Differences were considered significant if  $p < 0.05$ .

5           The results of the above studies are  
summarized in Table 3, below.

Table 3

	<u>Expression on Coronary Arteries</u>	<u>Inhibitor Peptide</u>	<u>Control Peptide</u>
10	MHC Class II	$\pm^c$	++
	T cells	$\pm$	++
15	Macrophages	$\pm$	+
	ICAM-1	$\pm$	+
	VCAM-1	$\pm$	+
20	Total Fibronectin	$\pm$	++
	<u>Vessel Intimal Thickening</u>		
25	Percent of Vessels <sup>a</sup>	35 <sup>d</sup>	88
	Severity (Percent of Vessel area) <sup>b</sup>	16 <sup>e</sup>	36
30			

35       <sup>a</sup> Baseline (rabbit own host heart) of incidence was  
12 percent and 10 percent, respectively, for inhibitor  
and control peptide groups.

<sup>b</sup> Baseline (rabbit own host heart) of severity was 12 percent and 12 percent, respectively, for inhibitor and control peptide groups.

<sup>c</sup> Scoring was: -, negative; ±, minimal; +, little; ++, moderately abundant; +++, very abundant.

<sup>d</sup> p<0.001.<sup>e</sup>  $p < 0.001$ .



- 71 -

As is seen from the above results, use of a contemplated inhibitor peptide greatly reduced the inflammation-induced damage observed in the allografted hearts. These damage reductions are particularly evident in the vessel intimal thickening results, but are also seen less directly in the results relating to expressed inflammatory markers shown by the MHC Class II antigen, the increased presence of T cells and macrophages, and the total fibronectin.

Example 7: In Vitro Porcine Allograft Model

A similar study was carried out in vitro using porcine coronary artery endothelial cells (EC; as are present in the IEL) and smooth muscle cells (SMC; as are present in the medial layer of the artery). The two cell types were cultured using a membrane transwell system, with the SMC on the bottom layer in M-199 medium (Gibco Labs.). The SMC were stimulated with 100 ng/ml of interleukin-1 $\beta$  (IL-1 $\beta$ ) for 24 hours prior to the start of the assay. Porcine peripheral blood lymphocytes were separated by Ficoll-Hypaque, radiolabeled and incubated overnight (about 18 hours) on the EC.

Transendothelial lymphocyte migration in the IL-1 $\beta$ -stimulated SMC was observed as compared to unstimulated SMC ( $p < 0.05$ ). The inhibitor peptide of Example 6, phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub>, present at 10  $\mu$ g/ml in the medium reduced lymphocyte migration by about 30 percent ( $p < 0.05$ ), whereas the same amount of a control peptide whose sequence was scrambled did not reduce migration.

Increased expression of EC and SMC fibronectin and IL-1 $\beta$  are features of an immunoinflammatory response associated with accelerated graft arteriopathy following piglet heterotopic cardiac transplantation. The above

- 72 -

results indicate that IL-1 $\beta$  induces fibronectin production in this in vitro model, which in turn contributes to transendothelial lymphocyte migration. The above results also illustrate that a contemplated inhibitor peptide can be used to reduce this immunoinflammatory response.

Example 8:        Experimental Autoimmune  
                  Encephalomyelitis in Mice

Experimental autoimmune encephalomyelitis (EAE) is a demyelinating disease of the central nervous system that can be induced in susceptible strains of mice and rats by immunization with myelin basic protein, proteolipid protein (PLP), or their immunodominant T cell determinants, or by injection of CD4-positive T cell clones specific for those determinants. EAE serves as an animal model of human multiple sclerosis. In both diseases, circulating leukocytes such as T cells and monocytes penetrate the blood/brain barrier and damage myelin, resulting in paralysis.

EAE was induced in female SJL/J mice (8 to 14 weeks old) by immunization on day zero with 50  $\mu$ g of a peptide corresponding to positions 139-151 of PLP emulsified in a 1:1 mixture of PBS and complete Freund's adjuvant (CFA). Each mouse was injected with 0.2 ml of the adjuvant emulsion subcutaneously (s.c.) at two sites in the hind flank. All mice received 10<sup>7</sup> killed Bordetella pertussis units in 100  $\mu$ l were injected intravenously 24 to 72 hours later.

Mice were observed daily, beginning at day 8 for clinical signs of EAE, and disease was scored on a scale of 0-5 as: 0 = no disease; 1 = floppy tail; 2 = moderate hind limb weakness; 3 = paraparesis; 4 = paraplegis with moderate forelimb weakness; 5 = quadriplegis or premoribund state.

- 73 -

The inhibitor peptide N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub> was administered intraperitoneally at 1 mg/mouse in 0.2 ml of incomplete Freund's adjuvant at days 8 and 9. A peptide having a scrambled sequence [N-phenylacetyl-Asp-Leu-Phe-D-Pro-NH<sub>2</sub>] was similarly administered to serve as a control. The relative potency of this control peptide is shown in Table 1 to be <1.

Summed or averaged scores for clinical signs were plotted vs. time. The area under the resulting curves was calculated between day 8 and day 35 to calculate percentage inhibition of EAE by an inhibitor peptide. The percent inhibition was calculated as follows:

$$\% \text{ Inhibition} = 100 - \frac{\text{Area of inhibitor peptide} + \text{control area}}{\text{Area of control peptide}} \times 100$$

Two exemplary plots through day 31 are shown in the graph of Fig. 6 in which the darkened circles are averaged scores for six mice treated with the inhibitor peptide and darkened squares are averaged scores for six mice that received the scrambled sequence control peptide. As can be seen, animals treated with an inhibitor peptide contemplated herein exhibited marked improvement in clinical signs as compared to those animals treated with the control peptide.

Example 9: CS-1 Expression in Human Rheumatoid Arthritis

Surgically-obtained synovial specimens from human rheumatoid arthritis (RA) patients were examined microscopically for the expression of the CS-1 peptide portion of fibronectin. Ultrathin sections of tissue were stained by the immunoperoxidase technique using anti-CS-1 antibodies, and were studied using

- 74 -

transmission electron microscopy. These studies showed that CS-1 was expressed on the luminal aspect of blood vessel endothelium, on the luminal plasma membrane. The plasma membrane of synoviocytes in the synovial intimal lining at the interface with the joint space was also stained. The CS-1 peptide portion was not found to be expressed in normal synovium.

Binding studies were carried out using the Jurkat T cell line and frozen RA synovial sections. Jurkat cell adhesion could be inhibited by anti-VLA-4 antibodies or the 10-mer CS-1 peptide portion (500  $\mu\text{g/ml}$ ) used as standard here (SEQ ID NO:3), but not with antibodies to VLA-5, VCAM-1-A or VCAM-1-B or a peptide in which the 10-mer sequence was scrambled. Stimulated MOLT-4 cells behaved similarly. These results are reported in Elices et al., J. Clin. Invest., 93:405-416 (January 1994).

A similar inhibition of binding of Jurkat cells to human RA synovial sections and not to normal synovial sections was observed using the inhibitor peptide N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub>. That peptide was used at its IC<sub>50</sub> value shown in Table 1 to be about 312 times less than the IC<sub>50</sub> value for the standard 10-mer. The absolute value of that IC<sub>50</sub> value is about 0.5  $\mu\text{molar}$ .

These results illustrate the importance of the CS-1 peptide portion and VLA-4 in a human chronic immunoinflammatory disease state, rheumatoid arthritis. These results also show that a contemplated inhibitor cell can inhibit the binding of inflammatory cells in this human immunoinflammatory disease state.

Example 10: Treatment of Asthmatic Sheep

Six asthmatic sheep were treated in a double blind cross-over study by Dr. William M. Abraham of the

- 75 -

Department of Research, Mount Sinai Medical Center, 4300 Alton Road, Miami Beach, FL 33140. The sheep had been shown to develop both early and late bronchial responses to inhaled Ascuris suum antigen. This study was carried out generally as described in Abraham et al., J. Clin. Invest., 93:776-787 (1994).

The inhibitor peptide used here was N-phenylacetyl-Leu-Asp-Phe-D-Pro-NH<sub>2</sub>, with a control peptide that comprised the previously described peptide having the same residues in a scrambled sequence. The vehicle for the nebulized peptides was phosphate-buffered saline. The peptides were administered at a dose of 1 mg/kg each, twice a day for three days prior to challenge, as well as 0.5 hours prior to and four hours post challenge on day 4. Because this was a crossover study, each animal received one or the other treatment, followed by a rest and then the other treatment. Each animal therefore served as its own control.

As can be seen from the graph of Fig. 7, animals treated with the inhibitor peptide exhibited less of a change from baseline specific lung resistance (SR<sub>L</sub>) on challenge and then more rapidly returned to their baseline pulmonary functions than did animals treated with the control peptide. In addition, pulmonary function remained at about baseline values from about 3-4 hours after challenge through the end of the study (8 hours post challenge), whereas the control peptide-treated animals had an increase in that pulmonary function (SR<sub>L</sub>).

Post challenge airway responsiveness was also assayed. Here, specific lung resistance, SR<sub>L</sub>, returned to baseline values 24 hours after challenge, however, the sheep were hyperresponsive to inhaled carbacol at that time. A comparison of PC<sub>400</sub> values on carbacol

- 76 -

inhalation prior to and 24 hours after challenge indicated that a much smaller dosage of carbachol [about 12 breath units (BU)] was required to increase the  $SR_L$  value for control peptide-treated animals 4-fold over a saline control value, as compared to the amount required for the inhibitor peptide-treated group (about 27 BU). The pre-challenge values here were about 20-23 BU, so that the inhibitor peptide caused the animals'  $SR_L$  to be greater than pre-challenge values, indicating a lessened response to carbachol than the pre-challenge response.

Although the present invention has now been described in terms of certain preferred embodiments, and exemplified with respect thereto, one skilled in the art will readily appreciate that various modifications, changes, omissions and substitutions may be made without departing from the spirit thereof.

- 77 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

- (A) NAME: Cytel Corporation
- (B) STREET: 3525 John Hopkins Court
- (C) CITY: San Diego
- (D) STATE: California
- (E) COUNTRY: United States of America
- (F) POSTAL CODE: 92121
- (G) TELEPHONE: 619-552-2794
- (H) TELEFAX: 619-552-8801

(ii) TITLE OF INVENTION: CS-1 PEPTIDOMIMETICS, COMPOSITIONS AND METHODS OF USING THE SAME

(iii) NUMBER OF SEQUENCES: 15

## (iv) COMPUTER READABLE FORM:

- (A) MEDIUM TYPE: Floppy disk
- (B) COMPUTER: IBM PC compatible
- (C) OPERATING SYSTEM: PC-DOS/MS-DOS
- (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

## (vi) CURRENT APPLICATION DATA:

- (A) APPLICATION NUMBER: PCT/US94/
- (B) FILING DATE: 5-DEC-1994

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Asp	Glu	Leu	Pro	Gln	Leu	Val	Thr	Leu	Pro	His	Pro	Asn	Leu	His	Gly
1				5				10						15	
Pro	Glu	Ile	Leu	Asp	Val	Pro	Ser	Thr							
			20					25							

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 12 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

INFORMATION FOR SEQ ID NO:3:

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 4 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

```
(ix) FEATURE:
      (A) NAME/KEY: Modified-site
      (B) LOCATION: 1
      (D) OTHER INFORMATION: /label= Xaa
                           /note= "Xaa= phenylacetyl-Leu."
```

```
(ix) FEATURE:
      (A) NAME/KEY: Modified-site
      (B) LOCATION: 4
      (D) OTHER INFORMATION: /label= Xaa
                             /note= "Xaa= Pro-NH2."
```

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 4 amino acids  
 (B) TYPE: amino acid  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide



- 79 -

## (ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = phenylacetyl-Leu."

## (ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 4
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = an amide formed between the  
C-terminal Pro carboxyl and a  
tetraethylenepentaamine containing 4  
N-phenylacetyl-Leu-Asp-Phe-Pro peptides  
bonded thereto."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Xaa Asp Phe Xaa  
1

## (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 4 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = GlcNAc-O-(CH<sub>2</sub>)<sub>5</sub>-C(=O)-Phe."

## (ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 4
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = Leu-piperidinamide."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Xaa Leu Asp Xaa  
1

## (2) INFORMATION FOR SEQ ID NO:7

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 4 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

- 80 -

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = benzoyl-Leu."

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 4
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = Pro-NH2."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Xaa Asp Val Xaa  
1

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = pivaloyl-Leu."

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 4
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = Pro-NH2."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Xaa Asp Val Xaa  
1

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 4 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- 81 -

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = phenylacetyl-Leu."

## (ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 4
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = Pro-decylamide."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Xaa Asp Phe Xaa  
1

## (2) INFORMATION FOR SEQ ID NO:10:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 10 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 10
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = Pro-NH2."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Ile Leu Asp Val Pro Ile Leu Asp Val Xaa  
1 5 10

## (2) INFORMATION FOR SEQ ID NO:11

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = Pro-NH2."

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Ile Leu Asp Phe Xaa  
1 5

- 82 -

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 4 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = phenylacetyl-Leu."

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 4
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = Pro-NH(CH2)5-C(O)NHC18H37."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Xaa Asp Phe Xaa  
1

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 5 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = Pro-NH2."

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Ile Leu Asp Val Xaa  
1 5

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 5 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

- 83 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Ile Leu Asp Val Pro  
1 5

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 1
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = acetyl-Ser."

(ix) FEATURE:

- (A) NAME/KEY: Modified-site
- (B) LOCATION: 5
- (D) OTHER INFORMATION: /label= Xaa  
/note= "Xaa = Ser-NH2."


(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Xaa Phe Asp Phe Xaa  
1 5

- 84 -

CLAIMS

1. A peptide of formula A



5

wherein B is an  $\alpha$ -hydrophobic amino acid residue whose side chain has a length of about 2 to about 5 methylene groups;

10 X is a group amide-linked to the nitrogen atom of B, said group having a ring structure bonded to the carbonyl carbon of said amide-linkage by a spacer having a length of zero to about two methylene groups, the length of X, including said spacer and carbonyl carbon, being about that of 3-quinolinecarbonyl or  
15 smaller, said ring structure being a 5- and 6-membered ring or a fused 6,6- or 6,5-membered ring; and

Z is selected from the group consisting of

(a) Xaa-NCy<sup>1</sup> where Xaa is Val, Ile, Leu  
20 or an amino acid residue having a side chain that contains one or two fused aromatic rings and NCy<sup>1</sup> is a cyclic ring-containing group having a ring nitrogen atom that forms an amide bond with the  $\alpha$ -carboxyl group of Xaa, and whose cyclic ring contains 5- or 6-atoms  
25 including said ring nitrogen atom; and

(b) NCy<sup>2</sup> where the depicted nitrogen is an amine substituent of a cyclic group whose depicted nitrogen atom forms an amide bond with the  $\alpha$ -carboxyl group of said Asp, and which amine substituent is bonded  
30 to a 6- or 7-membered ring or to a fused 6,6- or 6,7-membered lactam ring system in which the ring bearing said amine substituent is saturated and contains said amine substituent  $\alpha$  to the carbonyl group of said lactam;

- 85 -

said peptide inhibiting the binding of Jurkat cells (ATCC TIB 152) to a solid phase-bound peptide of SEQ ID NO:1 in an in vitro assay in an aqueous buffer at a pH value of 7.2-7.4 to an extent that is equal to or up to about 3000-fold greater than the inhibition in said binding exhibited by a peptide of SEQ ID NO:3.

2. The peptide according to claim 1 wherein B is selected from the group consisting of leucine, cyclohexylalanine, norleucine, methionine, homoserine, threonine, phenylalanine, valine, norvaline and isoleucine.

3. The peptide according to claim 2 wherein the ring structure of said X group contains an aromatic ring.


4. The peptide according to claim 3 wherein Z is Xaa-NCy<sup>1</sup>.

5. The peptide according to claim 4 wherein Xaa is an amino acid residue having a side chain that contains one or two fused aromatic rings.

6. The peptide according to claim 5 wherein the cyclic ring of NCy<sup>1</sup> is substituted with one or two substituent groups selected from the group consisting of carboxyl, C<sub>1</sub>-C<sub>4</sub> alkylencarboxyl, carboxamide, C<sub>1</sub>-C<sub>4</sub> alkylencarboxamide, hydroxyl, hydroxymethyl, (CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>H where n is 1, 2 or 3, and C<sub>1</sub>-C<sub>4</sub> alkyl.

- 86 -

7. A peptide of formula I



5                                wherein

                              X is a group amide-linked to the nitrogen  
atom of Leu, said group having a ring structure bonded  
to the carbonyl carbon of said amide-linkage by a spacer  
having a length of zero to about two methylene groups,  
10                               the length of X, including said spacer and carbonyl  
carbon, being about that of 3-quinolinecarbonyl or  
smaller, said ring structure being a 5- and 6-membered  
ring or a fused 6,6- or 6,5-membered ring; and

                              Z is selected from the group consisting  
15                               of

                              (a) Xaa-NCy<sup>1</sup> where Xaa is Val, Ile, Leu  
or an amino acid residue having a side chain that  
contains one or two fused aromatic rings and NCy<sup>1</sup> is a  
cyclic ring-containing group having a ring nitrogen atom  
20                               that forms an amide bond with the  $\alpha$ -carboxyl group of  
Xaa, and whose cyclic ring contains 5- or 6-atoms  
including said ring nitrogen atom; and

                              (b) NCy<sup>2</sup> where the depicted nitrogen is  
an amine substituent of a cyclic group whose depicted  
25                               nitrogen atom forms an amide bond with the  $\alpha$ -carboxyl  
group of said Asp, and which amine substituent is bonded  
to a 6- or 7-membered ring or to a fused 6,6- or  
6,7-membered lactam ring system in which the ring  
bearing said amine substituent is saturated and contains  
30                               said amine substituent  $\alpha$  to the carbonyl group of said  
lactam;

                              said peptide of formula I inhibiting the  
binding of Jurkat cells (ATCC TIB 152) to a solid phase-  
bound peptide of SEQ ID NO:1 in an in vitro assay in an  
35                               aqueous buffer at a pH value of 7.2-7.4 to an extent



- 87 -

that is about 10- to about 3000-fold better than the inhibition in said binding exhibited by a peptide of SEQ ID NO:3.

5                   8.    The peptide according to claim 7 wherein the ring structure of said X group contains an aromatic ring.

                  9.    The peptide according to claim 8 wherein  
10   said aromatic ring is substituted with a C<sub>1</sub>-C<sub>2</sub> alkyl or hydroxyl group.

                  10.   The peptide according to claim 8 wherein  
15   Z is Xaa-NCy<sup>1</sup>.

                  11.   The peptide according to claim 10 wherein  
Xaa is an amino acid residue having a side chain that contains one or two fused aromatic rings.

20                  12.   The peptide according to claim 7 wherein Z is Xaa-NCy<sup>1</sup>, and Xaa is an amino acid residue whose side chain has one or two fused aromatic rings.

                  13.   The peptide according to claim 12 wherein  
25   the cyclic ring of NCy<sup>1</sup> is substituted with one or two substituent groups selected from the group consisting of carboxyl, C<sub>1</sub>-C<sub>4</sub> alkylencarboxyl, carboxamide; C<sub>1</sub>-C<sub>4</sub> alkylencarboxamide, hydroxyl, hydroxymethyl, (CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>H where n is 1, 2 or 3, and C<sub>1</sub>-C<sub>4</sub> alkyl.

30                  14.   The peptide according to claim 7 wherein Z is NCy<sup>2</sup>.


- 88 -

15. The peptide according to claim 12 wherein NCy<sup>2</sup> is selected from the group consisting of benzylamido, phenethylamido, 2-(N-morpholinyl)ethylamido, N-carboxamidomethyl-N-(3-caprolactamyl)amido, N-(3-caprolactamyl)amido, N-(3-valerolactamyl)amido, N-2-(4-morpholinyl)ethyl-N-(2-oxo-tetrahydroquinolin-3-yl)amido, and N-(2-oxo-tetrahydroquinolin-3-yl)amido groups.

16. The peptide according to claim 7 wherein X is an amino acid residue having a cyclic ring side chain.

17. The peptide according to claim 16 wherein the nitrogen atom of the  $\alpha$ -amino group of the X amino acid residue is acylated with a C<sub>1</sub>-C<sub>6</sub> acyl group.

18. A peptide of the formula



wherein Ar is pyrazolyl, phenyl, pyridyl, or 3-quinolinyl;

Y is a spacer that is absent, -CH<sub>2</sub>-, -CH(NH)-, -O- or -NH-;

or Ar-Y-C(O)- together with the nitrogen atom of said Leu forms a phthalimido, a 1,2,3,4-tetrahydroquinazoline-2,4-dione-3-yl or 5-phenyldantoin-3-yl group, and

Ar-Y-C(O)- has a length of about 3-quinolinecarbonyl or less;

Xaa is an aromatic amino acid residue; and

NCy<sup>1</sup> is an amine-containing 5- or 6-membered cyclic ring group whose depicted nitrogen

- 89 -

atom is within the ring and forms an amide bond with  $\alpha$ -carboxyl of Xaa.

5           19.    The peptide according to claim 18 wherein Y is absent.

            20.    The peptide according to claim 18 wherein Xaa is selected from the group consisting of phenylalanyl, tryptophyl and tyrosyl.

10           21.    The peptide according to claim 18 wherein Ar-Y-C(O)- is a substituent selected from the group consisting of benzoyl, phenylacetyl, 3-pyridinecarbonyl, 4-pyridinecarbonyl, 3-pyridineacetyl, phenoxycarbonyl, 15    anilino carbonyl, pyrazolecarbonyl, tryptophyl and 3,4-dihydroxybenzoyl groups.


            22.    The peptide according to claim 18 wherein NCy<sup>1</sup> is selected from the group consisting of 20    morpholinamido, thiomorpholinamido, 4-(thiadioxo)piperidinamido, piperidamido, 4-substituted piperidinamido whose substituent is selected from the group consisting of hydroxy, carbamyl and carboxyl, L-2-(carboxamide)pyrrolidinamido, D-2-(carboxamide)- 25    pyrrolidinamido, pyrrolidinamido, 3,4-dihydroxy-pyrrolidinamido, 2-(hydroxymethyl)pyrrolidinamido, piperazinamido and 4-substituted piperazinamido whose substituent is selected from the group consisting of carboxymethyl, carboxamidomethyl, (5-hydroxyethylenoxy- 30    ethylene) and p-toluenesulfonamido groups.

            23.    The peptide according to claim 22 wherein Ar-Y-C(O)- together with the nitrogen atom of said Leu form a phthalimido group.

35


- 90 -

24. A peptide of the formula



5                    wherein NCy<sup>3</sup> is selected from the group  
consisting of morpholinamido, thiomorpholinamido,  
4-(thiadioxo)piperidinamido, D-2-(carboxamide)-  
pyrrolidinamido, piperidinamido, 4-substituted  
10                   piperidinamido whose substituent is selected from the  
group consisting of hydroxy, carbamyl and carboxyl,  
piperazinamido and 4-substituted piperazinamido whose  
substituent is selected from the group consisting of  
carboxymethyl, carboxamidomethyl, (5-hydroxyethylen-  
oxyethylene) and p-toluenesulfonamido and  
15                   pyrrolidinamido groups.

25. A process for treating fibronectin  
CS-1/VLA-4-mediated inflammation that comprises  
administering to a mammal having said inflammation an  
20                   inflammation-reducing amount of a peptide of the formula



                     wherein  
25                    X is a group amide-linked to the nitrogen  
atom of Leu, said group having a ring structure bonded  
to the carbonyl carbon of said amide-linkage by a spacer  
having a length of zero to about two methylene groups,  
the length of X, including said spacer and carbonyl  
30                   carbon, being about that of 3-quinolinecarbonyl or  
smaller, said ring structure being a 5- and 6-membered  
ring or a fused 6,6- or 6,5-membered ring; and  
                     Z is selected from the group consisting  
of

- 91 -

(a) Xaa-NCy<sup>1</sup> where Xaa is Val, Ile, Leu or an amino acid residue having a side chain that contains one or two fused aromatic rings and NCy<sup>1</sup> is a cyclic ring-containing group having a ring nitrogen atom that forms an amide bond with the  $\alpha$ -carboxyl group of Xaa, and whose cyclic ring contains 5- or 6-atoms including said ring nitrogen atom; and

(b) NCy<sup>2</sup> where the depicted nitrogen is an amine substituent of a cyclic group whose depicted nitrogen atom forms an amide bond with the  $\alpha$ -carboxyl group of said Asp, and which amine substituent is bonded to a 6- or 7-membered ring or to a fused 6,6- or 6,7-membered lactam ring system in which the ring bearing said amine substituent is saturated and contains said amine substituent  $\alpha$  to the carbonyl group of said lactam;

said peptide of formula I inhibiting the binding of Jurkat cells (ATCC TIB 152) to a solid phase-bound peptide of SEQ ID NO:1 in an in vitro assay in an aqueous buffer at a pH value of 7.2-7.4 to an extent that is about 10- to about 3000-fold better than the inhibition in said binding exhibited by a peptide of SEQ ID NO:3.

26. The process according to claim 25 wherein said CS-1-mediated inflammation is asthma.

27. The process according to claim 26 wherein said administration is by inhalation.

28. The process according to claim 25 wherein said administration is parenteral.

- 92 -

29. The process according to claim 25 wherein said CS-1-mediated inflammation is rheumatoid arthritis or osteoarthritis.

5           30. The process according to claim 25 wherein said CS-1-mediated inflammation is allograft rejection.

31. The process according to claim 25 wherein said CS-1-mediated inflammation is a skin inflammation.

10

32. The process according to claim 25 wherein said CS-1-mediated inflammation is a central nervous system demyelinating disease.

15

33. A pharmaceutical composition comprising an inflammation-reducing amount of a peptide dissolved or dispersed in a pharmaceutically acceptable diluent, said peptide having the formula

20

X-Leu-Asp-Z

wherein

25           X is a group amide-linked to the nitrogen atom of Leu, said group having a ring structure bonded to the carbonyl carbon of said amide-linkage by a spacer having a length of zero to about two methylene groups, the length of X, including said spacer and carbonyl carbon, being about that of 3-quinolinecarbonyl or smaller, said ring structure being a 5- and 6-membered ring or a fused 6,6- or 6,5-membered ring; and

30

Z is selected from the group consisting of

35           (a) Xaa-NCy<sup>1</sup> where Xaa is Val, Ile, Leu or an amino acid residue having a side chain that contains one or two fused aromatic rings and NCy<sup>1</sup> is a

- 93 -

cyclic ring-containing group having a ring nitrogen atom that forms an amide bond with the  $\alpha$ -carboxyl group of Xaa, and whose cyclic ring contains 5- or 6-atoms including said ring nitrogen atom; and

5 (b) NCy<sup>2</sup> where the depicted nitrogen is an amine substituent of a cyclic group whose depicted nitrogen atom forms an amide bond with the  $\alpha$ -carboxyl group of said Asp, and which amine substituent is bonded to a 6- or 7-membered ring or to a fused 6,6- or  
10 6,7-membered lactam ring system in which the ring bearing said amine substituent is saturated and contains said amine substituent  $\alpha$  to the carbonyl group of said lactam;

said peptide of formula I inhibiting the  
15 binding of Jurkat cells (ATCC TIB 152) to a solid phase-bound peptide of SEQ ID NO:1 in an in vitro assay in an aqueous buffer at a pH value of 7.2-7.4 to an extent that is about 10- to about 3000-fold better than the inhibition in said binding exhibited by a peptide of SEQ  
20 ID NO:3.

34. The pharmaceutical composition according to claim 33 wherein the ring structure of said X group contains an aromatic ring.

25


35. The pharmaceutical composition according to claim 33 wherein Z is Xaa-NCy<sup>1</sup>.

36. The pharmaceutical composition according to claim 33 wherein Z is Xaa-NCy<sup>1</sup>, and Xaa is a amino acid residue whose side chain has one or two fused aromatic rings.

30

- - 94 -

37. The pharmaceutical composition according to claim 36 wherein X has the formula



5

wherein Ar is pyrazolyl, phenyl, pyridyl, or 3-quinolinyl;

Y is a spacer that is absent,  $-\text{CH}_2-$ ,  $-\text{CH}(\text{NH})-$ ,  $-\text{O}-$  or  $-\text{NH}-$ ;

10

or  $\text{Ar-Y-C(O)-}$  together with the nitrogen atom of said Leu forms a phthalimido, a 1,2,3,4-tetrahydroquinazoline-2,4-dione-3-yl or 5-phenyldantoin-3-yl group, and

15


$\text{Ar-Y-C(O)-}$  has a length of about 3-quinolinecarbonyl or less.

38. The pharmaceutical composition according to claim 37 wherein  $\text{Ar-Y-C(O)-}$  is phenylacetyl, Xaa is Phe and  $\text{NCy}^1$  is  $\text{NCy}^3$  that is selected from the group consisting of morpholinamido, thiomorpholinamido, 4-(thiadioxo)piperidinamide, D-2-(carboxamide)-pyrrolidinamido, piperidinamido, 4-substituted piperidinamido whose substituent is selected from the group consisting of hydroxy, carbamyl and carboxyl, 25 piperazinamido and 4-substituted piperazinamido whose substituent is selected from the group consisting of carboxymethyl, carboxamidomethyl, (5-hydroxyethylenoxyethylene) and p-toluenesulfonamido and pyrrolidinamido groups.


30




1/6

**Fig. 1**INHIBITORY POTENCY  
(% OF STANDARD)

2/6



**Fig. 2**INHIBITORY POTENCY  
(% OF STANDARD)

3/6

**Fig. 3**INHIBITORY POTENCY  
(RELATIVE TO STANDARD)


4/6

**Fig. 4A**



**Fig. 4B**


5/6

**Fig. 5**EAR SWELLING  
(X 10<sup>-4</sup> INCHES)

AVERAGE SCORE  
(N=6)

6/6

Fig. 6



SR<sub>L</sub>  
(% CHANGE FROM BASELINE)




Fig. 7

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(6) : C07K 5/08, 5/10, 7/06; A61K 38/06, 38/07, 38/08

US CL : 530/330, 331; 514/17, 18

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/330, 331; 514/17, 18

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS ONLINE, MEDLINE

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A, P	US, A, 5,294,551 (FURCHT ET AL) 15 March 1994, see entire document.	1-38
A	US, A, 4,822,606 (SNYDERMAN ET AL) 18 April 1989, see entire document.	1-38

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

## \* Special categories of cited documents:

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*Z\* document member of the same patent family

Date of the actual completion of the international search

07 FEBRUARY 1995

Date of mailing of the international search report

06 MAR 1995

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

CAROL A. SALATA

Telephone No. (703) 308-0196

**THIS PAGE BLANK (USPTO)**